CUMULATIVE AUTHOR INDEX
Environmental Entomology
Volume 33, 2004

A

Abramson, C. I., 378
Acar, E. B., 832
Aldrich, J. R., 12
Ali, A., 1282
Alma, A., 1689
Andersen, P. C., 165
Anderson, P. L., 1109, 1116
Anduaga, S., 1306
Appel, A. G., 197
Asaro, C., 397
Augustin, S., 1584
Awadzi, K. D., 1351

B

Babah, M.A.O., 1577
Babin, R., 1458
Baez, I., 662
Baker, J., 671
Bancroft, J., 435
Barlow, N. D., 941
Barratt, B.I.P., 941
Barron, M. C., 941
Bartelt, R. J., 1217
Baum, K. A., 727
Belden, J., 1562
Bell, H. A., 949
Belote, R. T., 1609
Berger, P. H., 1207
Berhow, M., 1217
Berisford, C. W., 397
Beyo, J., 1458
Biggs, R. W., 921
Blackmer, J. L., 88, 1389
Bläske, V.-U., 54
Blom, P. E., 267
Blossey, B., 564
Boggs, C. L., 1020
Boiteau, G., 1421
Boivin, G., 362, 855
Bommireddy, P. L., 1549
Bonjour, E. L., 426, 1465
Booth, G. M., 832
Bosque-Pérez, N. A., 1207
Botero-Garcés, N., 310
Boucias, D. G., 54
Bourne, A., 997
Brandenburg, R. L., 1431
Brewer, G. J., 1037
Briano, J. A., 1436
Broce, A. B., 75
Brodbeck, B. V., 165
Brodeur, J., 1088
Brower, L. P., 249
Bruck, D. J., 1335
Bryant, V. M., Jr., 727
Buckley, F. G., 1266
Buckley, P. A., 1266
Bulka, K. C., 182
Bullas-Appleton, E. S., 1381
Burke, S., 221
Buschman, L. L., 487, 930
Butler, L., 418
Bybee, L. F., 138, 1232
Byerly-Murphy, K. F., 1242

C

Calvin, D. D., 1116
Cameron, C. A., 1178
Cameron, R. S., 397
Campbell, J. F., 75
Campbell, K., 138, 1232
Campobasso, G., 1673
Cañas, L. A., 88
Cannon, C. A., 815
Cano-Santana, Z., 297
Cantliffe, D. J., 1014
Caprio, M. A., 1223
Cardoso, F. F., 608
Carrière, Y., 921
Carrillo, M. A., 815
Caveney, S., 227
Chandler, L. D., 554
Chandler, M. A., 147
Chen, B., 155
Chen, L.-M., 1321
Chen, T.-Y., 1416
Cheng, D., 1253
Chu, C.-C., 1416
Clark, T. L., 681, 740, 1497
Cleary, D.F.R., 528
Coats, J., 1562
Cocquempot, C., 844
Coleman, R. J., 443
Coll, M., 1471
Colonnelli, E., 1673
Colpetzer, K., 990
Cortesero, A. M., 362
Costamagna, A. C., 128
Coulson, R. N., 727
Craig, T. P., 709
Cronin, J. T., 1052
Cuda, J. P., 1351
Cuperus, G. W., 426

D

Dahlsten, D. L., 1554
Dau, B., 1020
Dávalos, A., 564
Davis, A. K., 249
Dedej, S., 100
De Graaf, J., 1431
De Groot, P., 221, 844
Delaplane, K. S., 100
Demmon, A. S., 1523
Dennehy, T. J., 921
Dias, C.T.D.S., 1192
Difonzo, C. D., 35, 608
Dillard, H. R., 1593
Ding, H., 1207
Dively, G. P., 1116
Dixon, P. L., 405
Donaldson, J. R., 1505
Dormont, L., 370
Dorn, S., 81, 1155, 1410
Dosdall, L. M., 756
Drapeau, P., 1703
Duan, J. J., 275
Dubois, T., 62

E

Economou, L. P., 1, 1137
Edgerly, J. S., 907
Edwards, J. P., 949
Eigenbrode, S. D., 1207
Elias, S. P., 900
Eliopoulos, P. A., 1
Elkin, C. M., 1070
Elliott, N. C., 776
Ellis, J. D., Jr., 794
Ellsbury, M. M., 554
Elzen, P. J., 794
Emmen, D. A., 890
Epinat, G., 370
Erbilgin, N., 1554
Errouissi, F., 45
Evenden, M. L., 213

F

Fadamiro, H. Y., 1711
Farrar, R. R., Jr., 982
Fei, H., 242
Feng, H., 1253
Ferguson, C. M., 941
Ferrier, S. M., 700
Ferro, D. N., 787
Fielding, D. J., 1513
Fields, P., 671
Figueroa-Castro, D. M., 297
Filotas, M. J., 1127
Fischer, J., 1266
Fischer, S., 81
Fitzgerald, G., 1416
Fleischer, S. J., 267, 890
Flinn, P., 671
Fondren, K. M., 1652

Forschler, B. T., 1145
Fournier, V., 1088
Fox, T. B., 608
Frank, S. D., 1662, 1681
French, B. W., 554, 1497
Frouz, J., 1282
Frouzova, J., 1282
Fuller, B. W., 554
Funderburk, J. E., 662
Furniss, M. M., 21
Fuxa, J. R., 975

G

Gao, R., 435, 644
Garzón-Tiznado, J. A., 1242
Gates, M. W., 426, 1465
Gilbert, M., 1584
Gillard, C., 1381
Gillespie, R. G., 1296
Gillette, N. E., 206
Ginsberg, H. S., 1266
Glynn, C., 748
Goeriz, R. E., 1362
Gong, P., 1570
González-Megías, A., 188
Goolsby, J. A., 997
Goubault, M., 362
Goyer, R. A., 1477
Gozé, E., 1458
Grant, J. F., 1274
Greenberg, S. M., 443
Grosman, D. M., 397
Gross, J., 799
Grue, C. E., 872
Gryuntal, S., 1442
Guichard, S., 1584
Guo, Y., 1253
Gutierrez Quispe, M. R., 477
Gutowski, J. M., 844

H

Haack, R. A., 221
Habeck, D. H., 1351
Hagenbuch, K. L., 779
Hagler, J. R., 88
Hajek, A. E., 62, 1127
Haloti, S., 45
Hamilton, G. C., 1185
Hamm, J. J., 633
Han, Q., 1223
Hanks, L. M., 291
Hanlon, C. C., 138, 1232
Hansen, L. D., 832
Hansen, R. W., 147
Hao, S.-G., 1528
Hardie, J., 1081
Harding, C. L., 267
Harkins, K. R., 990
Harvey, J. A., 1488
Hausmann, C., 1410
Hawkins, B. A., 765
Head, G., 275
Hébert, C., 1703
Heinz, K. M., 282
Hellmich, R. L., 1109, 1116
Henderson, G., 242
Hendrix, D. L., 690
Henneberry, T. J., 690, 1416
Hentz, M., 546
Hepburn, R., 794
Herms, D. A., 748
Hern, A., 1155
Hibbard, B. E., 681, 740, 1102, 1497
Hicks, S. K., 779
Hilker, M., 799
Hillier, N. K., 405
Hodgson, E. W., 1037
Holman, M. S., 900
Horton, D. L., 174
Hou, X., 671
Hough-Goldstein, J., 990
Hough-Goldstein, J. A., 320, 577, 1632
Hower, A., 890
Hower, A. A., 119
Hu, X. P., 197
Hughes, J., 1155
Hurd, L. E., 182
Hutchison, W. D., 815
Hwang, S.-Y., 1321

I

Isaacs, R., 310
Isard, S. A., 650, 860
Ives, A. R., 1523

J

Jackai, L., 471
Janati-Idrissi, A., 45
Jayas, D. S., 1003
Jay-Robert, P., 45
Jensen, A., 275
Jensen, J. G., 1178
Jiafu, H., 62
Jian, F., 1003
Jiménez-Martínez, E. S., 1207
Jing, X.-H., 113
Joern, A., 1617
Johnson, J. M., 1632
Johnson, K., 1274
Johnson, M. W., 1088
Jones, A. M., 182
Jones, D. B., 776
Jones, J. B., 599

K

Kang, L., 113, 155, 1528
Kenis, M., 844
Kim, I.-K., 540
Kim, K. S., 457
Klein, R., 1274
Klepzig, K. D., 28
Koch, R. L., 815
Kontodimas, D. C., 1
Koona, P., 471
Kopper, B. J., 28
Krespi, L., 362
Kroschel, J., 477
Kumar, H., 1025
Kurota, H., 1163

L

Lacombe, E. H., 900
Lagnaoui, A., 477
Lan, Z., 174
Landis, D. A., 35, 128, 356, 608
Larson, D. J., 405
Lashomb, J. H., 1185
Latchininsky, A. V., 257
Lavoie, B., 1062
Lawson, A. B., 1554
Le Blanc, J.-P. R., 1421
Lee, H.-P., 540
Legaspi, B. C., Jr., 1401
Legaspi, J. C., 839, 1200, 1401
Lenssen, A. W., 450
Leonard, B. R., 1095
Lerin, J., 107
Leskey, T. C., 389
Lessio, F., 1689
Levine, E., 650, 860
Lewis, J. L., 1145
Lewis, L. C., 1109
Li, D., 1570
Li, G., 435, 644
Li, Z., 62
Lim, U. T., 1344
Lindroth, R. L., 1505
Liu, Z., 1570
Lobinske, R. J., 1282
Lockwood, J. A., 257
Lopes, J.R.S., 1192
Lopez, J. E., 1151
Losey, J. E., 535
Lubelczyk, C. B., 900
Luckman, B., 794
Lumaret, J.-P., 45, 370
Luna, M. G., 765
Lundgren, J. G., 958
Lykouressis, D. P., 499, 1137
Lysyk, T. J., 328, 823

M

Mabry, T. R., 650, 860
MacDonald, K. E., 227
MacDonald, L., 844
Macrae, I. V., 1037
Mallis, R. E., 182
Mantey, K. D., 506
Marchosky, R. J., Jr., 709
Martel, V., 855
Martin, C. G., 1351
Matadha, D., 1185
Matthews, H. J., 949
McCullough, D. G., 1652

McFaul, A. C., 1593
Mclaughlin, J. R., 213
McQuate, G. T., 1722
Meagher, R. L., 881
Medal, J. C., 1351
Meffert, L. M., 779
Meitner, C. J., 249
Messing, R. H., 1313
Miao, Z., 206
Mill, D. D., 832
Millar, J. G., 138, 1232
Miller, J. R., 35
Milonas, P. G., 513
Mitchell, J. P., 1447
Mitra, S., 1266
Mizell, R. F., III, 165
Moeser, J., 1169
Moon, R. D., 147
Moran, P. J., 346
Mowery, S. V., 75
Mulder, P. G., Jr., 378
Mullen, M. A., 75
Mullinix, B. G., Jr., 1371
Mussen, E. C., 1151

N

Nagoshi, R. N., 881
Nansen, C., 426, 1465
Naranjo, S. E., 1389
Natwick, E. T., 1416
Nault, B. A., 1593
Nault, L. R., 1192
Nava-Camberos, U., 1242
Nelson, H. J., 1523
Nenon, J.-P., 362
Neven, L. G., 506
Newton, A. S., Jr., 1644
Nibouche, S., 1458
Nott, R., 221
Nowak, J. T., 397
Nuessly, G., 546

O

Oberhauser, K. S., 1062
Oden, S., 165
Oi, D. H., 340
Oliveira, C.M.D., 1192
Olson, C., 921
O'Neal, M. E., 35
Opgenorth, D. C., 1644
Osisanya, E. O., 471
Otis, G., 1381
Oyediran, I. O., 740, 1497

P

Pacchioli, M. A., 119
Padgett, G. B., 1095
Paine, T. D., 138, 1232
Panizzi, A. R., 1217
Parajulee, M. N., 690, 1549
Parry, D., 1477
Payton, M. E., 426
Peck, S. L., 1722
Pedrosa-Macedo, J. H., 1351
Peng, C.Y.S., 1151
Perdikis, D. C., 499, 1137
Pereira, R. M., 340
Perez-Mendoza, J., 671, 807
Pesek, J. D., 320
Peterson, R.K.D., 450
Phillips, T. W., 426, 1465
Pinchak, W. E., 690
Poinsot, D., 362
Poland, T. M., 221
Porter, D. O., 1549
Porter, S. D., 1436
Potter, D. A., 520, 619
Prasifka, J. R., 282
Price, J. F., 1014
Price, P. W., 700
Prokopy, R. J., 1695

Q

Qureshi, J. A., 487, 930

R

Raffa, K. F., 28, 1554
Ragsdale, D. W., 147
Raimondo, S., 418
Ramaswamy, S. B., 487, 930
Rand, P. W., 900
Rao, A., 587, 1626
Razzak, A. A., 958
Reagan, T. E., 627
Reed, G., 275
Reid, M. L., 1070
Reitz, S. R., 662
Richter, A. R., 975
Ring, D. R., 1541
Rogers, M. E., 520, 619
Rondon, S. I., 1014
Rooks, E. C., 907
Rose, R., 1116
Rosenheim, J. A., 1088, 1362
Rowney, D. L., 1554
Ruberson, J. R., 964
Rubink, W. L., 727
Rull, J., 1695
Russo, J. M., 1116
Rutherford, B., 1274
Ryan, P. J., 1523
Rypstra, A. L., 1032

S

Saint-Germain, M., 1703
Salazar, J. A., 627
Samietz, J., 81, 1410
Sánchez-Piñero, F., 188
Sanders, N. J., 1609
Sandeson, P. D., 1421
Sansone, C. G., 282
Sappington, T. W., 443, 457
Savopoulou-Soultani, M., 513
Schaafsma, A. W., 1381
Scherm, H., 174
Schmolz, E., 799
Schoeman, A. S., 1431
Schultz, G., 1562
Schwerk, A., 1442
Sears, M. K., 1109, 1116
Sebolt, D. C., 356
Seckinger, J. O., 397
Sei, M., 1535
Sétamou, M., 443
Shah, D. A., 1593
Shannon, C. L., 450
Shapiro, M., 982
Shepard, B. M., 982
Sheridan, A., 378
Shrewsbury, P. M., 1662, 1681
Shufran, K. A., 776
Silva-Torres, C.S.A., 964
Simbro, E., 1562
Simmons, A. M., 839
Simmons, G. S., 88
Sisterson, M. S., 921
Six, D. L., 1554
Skinner, L. C., 147
Slosser, J. E., 690
Smith, B. N., 832
Smith, I., 949
Smith, J. W., Jr., 627
Smith, M. T., 435, 990
Smith, R. P., Jr., 900
Smith, S., 844
Snyder, W. E., 1523
Spencer, J. L., 650, 860
Spencer, N. R., 1673
Sporleder, M., 477
Squire, J., 378
Stanley-Horn, D. E., 1116
Stapleton, J. J., 1447
Stathas, G. J., 1
Stephens, E. J., 535
Stout, M. J., 720, 1541
Strazanac, J. S., 418
Styer, E. L., 633
Sumerford, D. V., 1109
Summers, C. G., 1447, 1644
Summerville, K. S., 234
Sun, J., 206
Svatoš, A., 1584
Sweeney, J., 844
Sword, G. A., 1577
Szyszko, J., 1442

T

Tabashnik, B. E., 921
Tamayo, M., 872
Taylor, G., 1274
Terragitti, G., 1673
Thacher, E. F., 1362
Throne, J. E., 487, 807, 930
Tillman, P. G., 633, 1371
Tindall, K. V., 720
Tobin, P. C., 435

Tonye, J., 471
Tooker, J. F., 291
Torres, J. B., 964
Tosh, C. R., 1081

U

Ulmer, B. J., 756
Underwood, H. B., 1266
Urías-López, M. A., 1242

V

Valles, S. M., 340
Vamvatsikos, P. G., 1081
Vance, R. A., 1274
Vandergast, A. G., 1296
Van Driesche, R. G., 1344
Vangessel, M. J., 320, 577, 1632
van Ginkel, W., 528
Vazquez, R. J., 1436
Vela-Coiffier, E. L., 1465
Velázquez-Monreal, J. J., 1242
Venette, R. C., 815
Verret, F., 1266
Vidal, S., 1169
Vinson, S. B., 587, 1602, 1626
Vogt, J. T., 1045, 1718

W

Wäckers, F. L., 81
Wagenaar, R., 1488
Wagner, D., 599
Wakarchuk, D., 221
Walker, K. R., 1327
Walter, A. J., 1652
Wang, B., 787
Wang, Q., 657
Wang, S., 787
Wang, X.-G., 1313
Weathersbee, A. A., III, 776
Weaver, D. K., 807
Weeks, R. D., Jr., 1602
Welter, S. C., 1327
Weltzin, J. F., 1609
West, M., 554
Whaley, P. M., 487, 930
White, N.D.G., 1003
White, W. H., 627
Wiedenmann, R. N., 958
Wiggins, G. J., 1274
Wilder, S. M., 1032
Williams, B. J., 720
Williams, D. W., 540, 644
Williams, L. H., III, 1389
Williams, M. A., 1052
Willrich, M. M., 1095
Wilson, A. P., 320, 577
Wilson, L. T., 1602
Wilson, T. A., 1102
Windham, M. T., 1274
Witjes, L.M.A., 1488
Wright, S. E., 389
Wu, J., 787
Wu, K., 1253, 1570

Y

Yuval, B., 1471

Z

Zemetra, R. S., 1207
Zeng, W., 657
Zhang, Q.-H., 12
Zhang, Z., 206, 206
Zhioua, E., 1266
Zhu, J., 1562
Zimmerman, K. M., 257
Zonneveld, R., 997
Zou, L., 1541

CUMULATIVE SUBJECT INDEX
Environmental Entomology
Volume 33, 2004

A

acclimation,
  cold hardiness, 155
  latitudinal variation, 155
  Liriomyza huidobrensis, 155
  pea leafminer, 155
Acer mono,
  Anoplophora glabripennis, 540
  Asian longhorned beetle, 540
  invasive species, 540
  South Korea, 540
acoustic communication,
  mole cricket, 1431
  moon phase, 1431
  sex ratio, 1431
Acyrthosiphon pisum, 1523, 1593
  aphid, 1523
  Aphidius ervi, 1523
  Aphis glycines, 1593
  host modification, 1523
  host–parasitoid interactions, 1523
  landscape ecology, 1593
  Rhopalosiphum maidis, 1593
  Therioaphis trifolii, 1593
adaptation,
  generalist, 1477
  oviposition, 1477
  polyphagy, 1477
  specialist, 1477
Adoxophyes orana,
  diapause termination, 513
  overwintering, 513
  photoperiod, 513
  Tortricidae, 513
Aesculus hippocastanum,
  biological invasion, 1584
  Cameraria ohridella, 1584
  pheromone trapping, 1584
  spatial dynamics, 1584
Aethina tumida,
  Apis mellifera, 794
  pupation success, 794
  soil properties, 794
Ageniaspis fuscicollis,
  field sex ratio, 528
  The Netherlands, 528
  Yponomeuta, 528
aggregation,
  gregarization, 1577
  phase, 1577
  Schistocerca gregaria, 1577
alate,
  apterous, 546
  Sipha elegans, 546
  yellow sugarcane aphid, 546
alfalfa, 119, 450, 890
  clover root curculio, 119
  Empoasca fabae, 890
  gas exchange, 450
  herbivory, 450
  Medicago sativa, 119, 890
  Medicago spp., 450
  nodules, 119
  proportion maps, 890
  red clover, 450
  Sitona hispidulus, 119
  temporal spatial dynamics, 890
alternate hosts, 681, 740, 1102, 1497
  Bacillus thuringiensis, 1102
  Diabrotica barberi, 1497
  Diabrotica virgifera virgifera, 681, 740, 1102
  grassy weeds, 1497
  host-searching behavior, 1497
  northern corn rootworm, 1497
  prairie grasses, 740
  resistance management, 681, 740, 1102
  rootworm, 681
  transgenic-rootworm maize, 1102
amino acids,
  dietary nitrogen, 165
  glassy-winged sharpshooter, 165
  Homalodisca coagulata, 165
  insect rearing, 165
amorphogenic effects,
  Apis mellifera, 1151
  fungicides, 1151
  honey bee, 1151
  toxic effects, 1151
anaerobic protists,
  physiological saline solution, 1145
  protist counting techniques, 1145
  Reticulitermes flavipes, 1145
  symbiotic protists, 1145
anautogenous,
  grass flies, 12
  hexyl butanoate, 12
  kairomone, 12
  Olcella, 12
Anisopteromalus calandrae,
  Cephalonomia waterstoni, 671
  Cryptolestes, 671
  Pea protein, 671
  Sitophilus, 671
Anoplophora glabripennis, 435, 540, 644
  Acer mono, 540
  Asian longhorned beetle, 540, 644
  dispersal, 644
  eradication, 435
  harmonic radar, 644
  individual mark–recapture, 644
  invasive species, 435, 540
  mark–recapture, 435
  South Korea, 540
  spatial autocorrelation, 435
Anthonomus grandis,
  gene flow, 457
  genetic diversity, 457
  mitochondrial DNA restriction fragment length polymorphism, 457
  phylogeography, 457
Anthonomus grandis grandis,
  boll weevil, 443
  cotton fruit sizes, 443
  feeding punctures, 443
  oviposition punctures, 443
Anthonomus pomorum,
  colors, 1410
  host finding, 1410
  host plant, 1410
  vision, 1410
antipredator behavior,
  biological control, 1032
  introduced predator, 1032
  Pardosa milvina, 1032
  Tenodera aridifolia sinensis, 1032
aphid, 776, 1523
  Acyrthosiphon pisum, 1523
  Aphidius ervi, 1523
  biological control, 776
  cold tolerance, 776
  geographic variation, 776
  host modification, 1523
  host–parasitoid interactions, 1523
  parasitoid, 776
aphid-borne viruses,
  Aphis gossypii, 1447
  Bemisia argentifolii, 1447
  spectral quantum flux, 1447
  UV reflective mulch, 1447
  wheat straw mulch, 1447

Aphidius ervi,
Acyrthosiphon pisum, 1523
aphid, 1523
host modification, 1523
host–parasitoid interactions, 1523
aphids, 1014, 1242
Bemisia tabaci, 1242
Coleomegilla, 1014
Geocoris, 1014
Orius, 1014
papaya ring spot virus, 1242
predators, 1014
squash leaf curl virus, 1242
twospotted spider mite, 1014
watermelon, 1242
Aphis fabae,
Cirsium arvense, 1081
clones, 1081
hybridization, 1081
plant species preference, 1081
Aphis glycines,
Acyrthosiphon pisum, 1593
landscape ecology, 1593
Rhopalosiphum maidis, 1593
Therioaphis trifolii, 1593
Aphis gossypii, 690, 1447
aphid-borne viruses, 1447
Bemisia argentifolii, 1447
cotton, 690
cotton aphid, 690
glucose, 690
leaf sugars, 690
spectral quantum flux, 1447
sucrose, 690
UV reflective mulch, 1447
wheat straw mulch, 1447
Aphthona lacertosa,
Aphthona nigriscutis, 147
biological control, 147
developmental rates, 147
Euphorbia esula, 147
leafy spurge, 147
Aphthona nigriscutis,
Aphthona lacertosa, 147
biological control, 147
developmental rates, 147
Euphorbia esula, 147
leafy spurge, 147
Apis mellifera, 100, 727, 794, 1151
Aethina tumida, 794
amorphogenic effects, 1151
coastal prairie, 727
feral colony, 727
fungicides, 1151
honey bee, 727, 1151
nectar robbing, 100
pollen, 727
pupation success, 794
rabbiteye blueberry, 100
soil properties, 794
toxic effects, 1151
Vaccinium ashei, 100
Xylocopa virginica, 100
apple maggot,
genotype, 1695
host finding, 1695
ovipositional boring, 1695
preference, 1695
apterous,
alate, 546
Sipha elegans, 546
yellow sugarcane aphid, 546
aquatic weed,
damage, 346
plant quality, 346
Pontederiaceae, 346
Argyrotaenia velutinana,
codling moth, 1711
Cydia pomonella, 1711
pest monitoring, 1711
redbanded leafroller, 1711
artificial substrate,
oviposition, 1217
plant extracts, 1217
Asclepias syriaca,
compensatory feeding, 1062
Danaus plexippus, 1062
Monarch butterflies, 1062
plant nitrogen, 1062
ascovirus transmission,
Campoletis sonorensis, 633
Cardiochiles nigriceps, 633
Heliothis virescens, 633
Microplitis croceipes, 633
Asian longhorned beetle, 62, 540, 644
Acer mono, 540
Anoplophora glabripennis, 540, 644
Beauveria bassiana, 62
Beauveria brongniartii, 62
dispersal, 644
entomopathogenic fungi, 62
harmonic radar, 644
individual mark–recapture, 644
invasive species, 540
Metarhizium anisopliae, 62
South Korea, 540
Asphondylia atriplicis,
Atriplex canescens, 709
Cecidomyiidae, 709
gall, 709
parasitoids, 709
assemblage composition,
dung beetle, 45
Scarabaeoidea, 45
spatial variability, 45
trophic resource, 45
Asteraceae, 291, 297
floral visitors, 297
parasitoid, 291
prairie, 291
predation, 291
Silphium, 291
similarity indexes, 297
stable isotope, 291
temporal variation, 297
xeric community, 297
Atriplex canescens,
Asphondylia atriplicis, 709
Cecidomyiidae, 709
gall, 709
parasitoids, 709
attract kill,
attracticide, 213
Grapholita molesta, 213
Oriental fruit moth, 213
pheromone, 213
attractants,
Cerambycidae, 844
host volatiles, 844
Tetropium fuscum, 844
traps, 844
attracticide,
attract kill, 213
Grapholita molesta, 213
Oriental fruit moth, 213
pheromone, 213
augmentative biological control,
balsam fir, 1652
Christmas trees, 1652
Chrysoperla rufilabris, 1652
green lacewings, 1652

B

Bacillus thuringiensis,
alternate hosts, 1102
Bacillus thuringiensis, 921, 982, 1335
Beauveria bassiana, 1335
Bt cotton, 921
corn, 982
Bacillus thuringiensis,
Diabrotica virgifera virgifera, 1102
Bacillus thuringiensis, 921, 982, 1025, 1335
diamondback moth, 1025
fall armyworm, 982
insect behavior, 1025
Metarhizium anisopliae, 1335
nontarget, 921
nucleopolyhedrovirus, 982
pink bollworm, 921
Plutella xylostella, 1025
Bacillus thuringiensis,
resistance management, 1102
Bacillus thuringiensis, 982, 1025, 1335
soil survey, 1335
Steinernema oregonense, 1335
transgenic crop, 1025
transgenic plants, 982

Bacillus thuringiensis,
transgenic-rootworm maize, 1102
balsam fir,
augmentative biological control, 1652
Christmas trees, 1652
Chrysoperla rufilabris, 1652
green lacewings, 1652
bark beetles,
insect–fungal interaction, 28
Ophiostoma ips, 28
ophiostomatoid fungi, 28
symbiosis, 28
Beauveria bassiana, 62, 1335
Asian longhorned beetle, 62
Bacillus thuringiensis, 1335
Beauveria brongniartii, 62
entomopathogenic fungi, 62
Metarhizium anisopliae, 62, 1335
soil survey, 1335
Steinernema oregonense, 1335
Beauveria brongniartii,
Asian longhorned beetle, 62
Beauveria bassiana, 62
entomopathogenic fungi, 62
Metarhizium anisopliae, 62
beech bark disease,
beech scale, 1274
Cryptococcus fagisuga, 1274
epidemiology, 1274
Nectria, 1274
beech scale,
beech bark disease, 1274
Cryptococcus fagisuga, 1274
epidemiology, 1274
Nectria, 1274
beetle,
biological control, 1673
Coleoptera, 1673
Thamnurgus euphorbiae, 1673
weed, 1673
behavior, 362, 860
crop rotation resistance, 860
Diabrotica virgifera virgifera, 860
external/internal marking, 362
feeding, 860
host quality, 362
oviposition, 860
superparasitism, 362
Bemisia argentifolii,
aphid-borne viruses, 1447
Aphis gossypii, 1447
spectral quantum flux, 1447
UV reflective mulch, 1447
wheat straw mulch, 1447
Bemisia tabaci, 839, 1242
aphids, 1242
biological control, 839
Delphastus catalinae, 839
papaya ring spot virus, 1242
squash leaf curl virus, 1242
vegetable, 839
watermelon, 1242
whitefly, 839
benzaldehyde,
Conotrachelus nenuphar, 389
grandisoic acid, 389
monitoring traps, 389
plum curculio, 389
big-eyed bug,
generalist predator, 964
Helicoverpa zea, 964
life table, 964
Spodoptera exigua, 964
biocontrol, 872, 1436
development time, 872
Euhrychiopsis lecontei, 872
host range, 1436
Solenopsis geminata, 1436
Solenopsis invicta, 1436
Solenopsis xyloni, 1436
watermilfoil, 872
biological control, 128, 147, 282, 340, 356, 499, 520, 587, 608, 662, 765, 776, 823, 839, 958, 1032, 1127, 1313, 1554, 1673, 1681
antipredator behavior, 1032
aphid, 776
Aphthona lacertosa, 147
Aphthona nigriscutis, 147
beetle, 1673
Bemisia tabaci, 839
Carabidae, 1681
chemical ecology, 1554
cold storage, 823
cold tolerance, 776
Coleoptera, 1673
conservation, 282
Delphastus catalinae, 839
developmental rates, 147
ectoparasitoids, 1313
Enoclerus lecontei, 1554
Enoclerus sphegeus, 1554
entomopathogenic fungi, 1127
Euphorbia esula, 147
fecundity, 128
fire ant, 340
Forelius, 587
fruit fly parasitoids, 1313
Galerucella calmariensis, 356
generalist predators, 662
geographic variation, 776
Hippodamia convergens, 282
host selection, 520
house fly, 823
hyperparasitism, 1313
inbreeding depression, 765
introduced predator, 1032
invasive species, 608
leafy spurge, 147
life history parameters, 128
longevity, 128
Lythrum salicaria, 356
Macrolophus pygmaeus, 499
Malacosoma disstria, 1127
mating status, 128
Miridae, 499
Monomorium, 587
Myzus persicae, 499
Nasonia vitripennis, 765
nontarget impact, 1313
outbreeding depression, 765
parasitoid, 520, 765, 776
Pardosa milvina, 1032
pathogen, 340
pepper, 499
Pheidole, 587
pollinivory, 958
polygyny, 340
predation, 356
predator community, 608
predator-prey interactions, 662
prey vulnerability, 662
pteromalids, 823
risk assessment, 958
S. molesta, 587
Scarabaeidae, 520
Scymnus loewii, 282
soil moisture, 1127
Solenopsis invicta, 340
Staphylinidae, 1681
survival, 823
Temnochila chlorodia, 1554
temperature, 1127
Tenodera aridifolia sinensis, 1032
Thamnurgus euphorbiae, 1673
transgenic crops, 958
turfgrass, 520, 1681
vegetable, 839
voracity, 1681
weed, 1673
whitefly, 839
Zea mays, 958
biological control weeds,
Florida Everglades restoration, 997
predictive studies, 997
biological invasion,
Aesculus hippocastanum, 1584
Cameraria ohridella, 1584
pheromone trapping, 1584
spatial dynamics, 1584
biotechnology,
Cry3Aa, 275
Leptinotarsa decemlineata, 275
nontarget effects, 275
potato pest management, 275
bison grazing,
Konza prairie, 1617
Orthoptera, 1617
population dynamics, 1617

prescribed burning, 1617
body size,
male aggression, 657
mating success, 657
Nadezhdiella cantori, 657
sexual selection, 657
boll rot,
cotton, 1095
hard-locked, 1095
stink bugs, 1095
boll weevil,
Anthonomus grandis grandis, 443
cotton fruit sizes, 443
feeding punctures, 443
oviposition punctures, 443
boll-feeding Lepidoptera,
sampling, 1458
within-plant distribution, 1458
boreal forest,
forest fire, 1703
habitat location, 1703
host selection, 1703
Monochamus scutellatus, 1703
bruchid,
diapause, 1163
overwintering stage, 1163
seasonal adaptation, 1163
Bt corn,
chronic effects, 1116
monarch butterfly, 1116
nontarget, 1116
risk assessment, 1116
Bt cotton,
Bacillus thuringiensis, 921
nontarget, 921
pink bollworm, 921
Bt maize,
dispersal, 487
elemental marking, 487
southwestern corn borer, 487

C

calorimetry,
Chrysomela lapponica, 799
color morph, 799
temperature, 799
thermal balance, 799
Calpodes ethlius,
ectoparasitoid, 227
Elachertus scutellatus, 227
koinobiont, 227
synovigenic, 227
Cameraria ohridella,
Aesculus hippocastanum, 1584
biological invasion, 1584
pheromone trapping, 1584
spatial dynamics, 1584
Campoletis sonorensis,
ascovirus transmission, 633
Cardiochiles nigriceps, 633
Heliothis virescens, 633
Microplitis croceipes, 633
canola,
host plant, 756
larval development, 756
polyphagous, 756
canonical correspondence analysis,
crop rotation, 554
Cry1Ab, 554
nontarget, 554
predators, 554
canopy temperature,
defoliation, 471
pod position, 471
Vigna unguiculata, 471
Carabidae, 564, 1442, 1681
biological control, 1681
feeding state, 1442
forest, 1442
garlic mustard, 564
geostatistics, 564
impact, 564
invasive, 564
male-to-female ratio, 1442
movement patterns, 1442
radar tracking, 1442
Staphylinidae, 1681
turfgrass, 1681
voracity, 1681
Cardiochiles nigriceps,
ascovirus transmission, 633
Campoletis sonorensis, 633
Heliothis virescens, 633
Microplitis croceipes, 633
caterpillar ecology,
Coenonympha tullia nipisiquit, 1535
hypoxia, 1535
salt marsh, 1535
tidal submergence, 1535
catnip,
elemol, 1562
nepetalactone, 1562
osage orange, 1562
repellency, 1562
cattle dung,
dung beetles, 370
horse dung, 370
olfactometer, 370
trophic preferences, 370
Cecidomyiidae,
Asphondylia atriplicis, 709
Atriplex canescens, 709
gall, 709
parasitoids, 709
Cephalonomia waterstoni,
Anisopteromalus calandrae, 671
Cryptolestes, 671
Pea protein, 671
Sitophilus, 671
Cerambycidae, 138, 844
attractants, 844
fecundity, 138
host volatiles, 844
longevity, 138
Phoracantha recurva, 138
Phoracantha semipunctata, 138
Tetropium fuscum, 844
traps, 844
chemical ecology,
biological control, 1554
Enoclerus lecontei, 1554
Enoclerus sphegeus, 1554
Temnochila chlorodia, 1554
Christmas berry,
herbivory, 1351
leafroller, 1351
life history, 1351
weed biological control, 1351
Christmas trees,
augmentative biological control, 1652
balsam fir, 1652
Chrysoperla rufilabris, 1652
green lacewings, 1652
chromatic contrast,
host location, 81
pupal parasitoid, 81
vision, 81
wavelength discrimination, 81
chronic effects,
Bt corn, 1116
monarch butterfly, 1116
nontarget, 1116
risk assessment, 1116
Chrysomela lapponica,
calorimetry, 799
color morph, 799
temperature, 799
thermal balance, 799
Chrysoperla rufilabris,
augmentative biological control, 1652
balsam fir, 1652
Christmas trees, 1652
green lacewings, 1652
Cirsium arvense,
Aphis fabae, 1081
clones, 1081
hybridization, 1081
plant species preference, 1081
clones,
Aphis fabae, 1081
Cirsium arvense, 1081
hybridization, 1081
plant species preference, 1081
clover root curculio,
alfalfa, 119
Medicago sativa, 119
nodules, 119
Sitona hispidulus, 119

CO potato beetle,
colonization, 267
immigration, 267
Leptinotarsa decemlineata, 267
mapping, 267
CO_2 enrichment,
global change, 1609
insect communities, 1609
invasive species, 1609
plant communities, 1609
coastal prairie,
Apis mellifera, 727
feral colony, 727
honey bee, 727
pollen, 727
coat protein–mediated resistance,
host plant resistance, 1207
insect behavior, 1207
virus vectors, 1207
virus-induced volatiles, 1207
Coccinellidae,
sweep-net sampling, 535
visual sampling, 535
yellow sticky cards, 535
codling moth,
Argyrotaenia velutinana, 1711
Cydia pomonella, 1711
pest monitoring, 1711
redbanded leafroller, 1711
Coenonympha tullia nipisiquit,
caterpillar ecology, 1535
hypoxia, 1535
salt marsh, 1535
tidal submergence, 1535
coevolution,
deme formation, 748
host plant resistance, 748
host specialization, 748
local adaptation, 748
cold hardiness, 113, 155, 815
acclimation, 155
fall generation, 113
Harmonia axyridis, 815
latitudinal variation, 155
Liriomyza huidobrensis, 155
Locusta migratoria, 113
low temperature, 815
overwintering, 815
pea leafminer, 155
postwinter population, 113
summer generation, 113
supercooling point, 815
cold storage,
biological control, 823
house fly, 823
pteromalids, 823
survival, 823
cold tolerance,
aphid, 776
biological control, 776
geographic variation, 776
parasitoid, 776
Coleomegilla,
aphids, 1014
Geocoris, 1014
Orius, 1014
predators, 1014
twospotted spider mite, 1014
Coleoptera,
beetle, 1673
biological control, 1673
Thamnurgus euphorbiae, 1673
weed, 1673
colonization,
CO potato beetle, 267
immigration, 267
Leptinotarsa decemlineata, 267
mapping, 267
color morph,
calorimetry, 799
Chrysomela lapponica, 799
temperature, 799
thermal balance, 799
colors,
Anthonomus pomorum, 1410
host finding, 1410
host plant, 1410
vision, 1410
compensatory feeding,
Asclepias syriaca, 1062
Danaus plexippus, 1062
Monarch butterflies, 1062
plant nitrogen, 1062
competition,
diapause, 1232
Phoracantha recurva, 1232
Phoracantha semipunctata, 1232
seasonal development, 1232
Conotrachelus nenuphar, 174, 389
benzaldehyde, 389
degree-day model, 174
grandisoic acid, 389
monitoring traps, 389
peach, 174
plum curculio, 174, 389
Prunus persica, 174
conservation,
biological control, 282
Hippodamia convergens, 282
Scymnus loewii, 282
conservation biological control, 619, 1662
habitat manipulation, 1662
supplemental food sources, 619
Tiphia pygidialis, 619
Tiphia vernalis, 619
turf, 1662
white grubs, 619
Coptotermes formosanus,
habitat temperature, 197
Reticulitermes flavipes, 197
seasonal variation, 197
thermal tolerances, 197
corn, 35, 807, 930, 982
Bacillus thuringiensis, 982
elemental marking, 930
fall armyworm, 982
life history, 807
modeling, 807
nucleopolyhedrovirus, 982
phenology, 35
resistance, 35
resistance management, 930
rotation, 35
Sitotroga cerealella, 807
southwestern corn borer, 930
soybean, 35
survivorship, 807
transgenic plants, 982
Zea mays, 930
corn earworm,
cotton, 1371
grain sorghum, 1371
Helicoverpa zea, 1371
trap crop, 1371
corn leafhopper,
corn stunt, 1644
Dalbulus maidis, 1644
spiroplasma, 1644
Spiroplasma kunkelii, 1644
corn stunt,
corn leafhopper, 1644
Dalbulus maidis, 1644
spiroplasma, 1644
Spiroplasma kunkelii, 1644
costs of mating,
longevity, 779
repeatability, 779
sexual selection, 779
starvation resistance, 779
Cotesia flavipes,
Diatraea saccharalis, 627
refuges, 627
sentinel larvae, 627
sugarcane, 627
cotton, 690, 1095, 1362, 1371
Aphis gossypii, 690
boll rot, 1095
corn earworm, 1371
cotton aphid, 690
diet selection, 1362
direct observations, 1362
glucose, 690
grain sorghum, 1371
habitat selection, 1362
hard-locked, 1095
Helicoverpa zea, 1371
leaf sugars, 690
omnivory, 1362
stink bugs, 1095
sucrose, 690
trap crop, 1371

cotton aphid,
Aphis gossypii, 690
cotton, 690
glucose, 690
leaf sugars, 690
sucrose, 690
cotton fruit sizes,
Anthonomus grandis grandis, 443
boll weevil, 443
feeding punctures, 443
oviposition punctures, 443
crop rotation,
canonical correspondence analysis, 554
Cry1Ab, 554
nontarget, 554
predators, 554
crop rotation resistance,
behavior, 860
Diabrotica virgifera virgifera, 860
feeding, 860
oviposition, 860
Cry1Ab,
canonical correspondence analysis, 554
crop rotation, 554
nontarget, 554
predators, 554
Cry3Aa,
biotechnology, 275
Leptinotarsa decemlineata, 275
nontarget effects, 275
potato pest management, 275
Cryptococcus fagisuga,
beech bark disease, 1274
beech scale, 1274
epidemiology, 1274
Nectria, 1274
Cryptolestes,
Anisopteromalus calandrae, 671
Cephalonomia waterstoni, 671
Pea protein, 671
Sitophilus, 671
Cryptolestes ferrugineus,
distribution, 1003
movement, 1003
stored grain, 1003
temperature gradient, 1003
wandering, 1003
Cyclosa,
edge effects, 1296
Hawaii, 1296
lava flows, 1296
Tetragnatha, 1296
Cydia (Grapholita) molesta,
dispersal, 1155
photoperiod, 1155
preimaginal environment, 1155
temperature, 1155
Cydia pomonella,
Argyrotaenia velutinana, 1711
codling moth, 1711
pest monitoring, 1711
redbanded leafroller, 1711
Cytospora,
feltleaf willow, 21
Scolytidae, 21
Trypophloeus, 21

D

Dalbulus maidis,
corn leafhopper, 1644
corn stunt, 1644
spiroplasma, 1644
Spiroplasma kunkelii, 1644
damage,
aquatic weed, 346
plant quality, 346
Pontederiaceae, 346
damage prediction,
pheromone traps, 397
Pinus taeda, 397
Rhyacionia frustrana, 397
Danaus plexippus, 1062, 1109
Asclepias syriaca, 1062
compensatory feeding, 1062
Monarch butterflies, 1062
nontargets, 1109
plant nitrogen, 1062
risk assessment, 1109
transgenic corn, 1109
decision support tools,
modeling, 426
Oklahoma, 426
stored-product beetles, 426
defoliation,
canopy temperature, 471
pod position, 471
Vigna unguiculata, 471
degree-day,
emergence, 1541
flight activity, 1541
Lissorhoptrus oryzophilus, 1541
Oryza sativa, 1541
degree-day model,
Conotrachelus nenuphar, 174
peach, 174
plum curculio, 174
Prunus persica, 174
Delphastus catalinae,
Bemisia tabaci, 839
biological control, 839
vegetable, 839
whitefly, 839
deme extinction,
introduced species, 182
mantids, 182
metapopulations, 182
Tenodera aridifolia sinensis, 182
deme formation,
coevolution, 748
host plant resistance, 748
host specialization, 748
local adaptation, 748
Dendroctonus ponderosae,
fire damage, 1070
habitat selection, 1070
population density, 1070
density,
dispersal, 1421
Leptinotarsa decemlineata, 1421
mark-release-recapture, 1421
detritivore,
insect ecology, 907
ordination, 907
webspinners, 907
development,
model, 1
Nephus bisignatus, 1
Nephus includens, 1
threshold, 1
development time,
biocontrol, 872
Euhrychiopsis lecontei, 872
watermilfoil, 872
developmental duration,
Lygus elisus, 1549
population dynamics, 1549
temperature, 1549
developmental polymorphism,
grass webworm, 1178
head capsule widths, 1178
developmental rate,
Encarsia citrina, 1185
euonymus scale, 1185
life table, 1185
Quadraspidiotus perniciosus, 1185
developmental rates,
Aphthona lacertosa, 147
Aphthona nigriscutis, 147
biological control, 147
Euphorbia esula, 147
leafy spurge, 147
Diabrotica barberi,
alternate hosts, 1497
grassy weeds, 1497
host-searching behavior, 1497
northern corn rootworm, 1497
Diabrotica virgifera virgifera, 650, 681, 740, 860, 1102
alternate hosts, 681, 740, 1102
Bacillus thuringiensis, 1102
behavior, 860
crop rotation resistance, 860
diel periodicity, 650
dispersal, 650
feeding, 860
flight, 650
oviposition, 860

prairie grasses, 740
resistance management, 681, 740, 1102
rootworm, 681
transgenic-rootworm maize, 1102
western corn rootworm, 650
diamondback moth,
Bacillus thuringiensis, 1025
insect behavior, 1025
Plutella xylostella, 1025
transgenic crop, 1025
diapause, 1163, 1232
bruchid, 1163
competition, 1232
overwintering stage, 1163
Phoracantha recurva, 1232
Phoracantha semipunctata, 1232
seasonal adaptation, 1163
seasonal development, 1232
diapause termination,
Adoxophyes orana, 513
overwintering, 513
photoperiod, 513
Tortricidae, 513
Diatraea saccharalis,
Cotesia flavipes, 627
refuges, 627
sentinel larvae, 627
sugarcane, 627
diel periodicity,
Diabrotica virgifera virgifera, 650
dispersal, 650
flight, 650
western corn rootworm, 650
diet selection,
cotton, 1362
direct observations, 1362
habitat selection, 1362
omnivory, 1362
dietary nitrogen,
amino acids, 165
glassy-winged sharpshooter, 165
Homalodisca coagulata, 165
insect rearing, 165
diffusion model,
glassy-winged sharpshooter, 88
IgG protein markers, 88
smoke tree sharpshooter, 88
yellow sticky traps, 88
Diplolepis spp.,
plant-stress hypothesis, 1052
plant-vigor hypothesis, 1052
Rosa arkansana, 1052
tall-grass prairie, 1052
Diptera,
environmental stress, 1282
flooding, 1282
Isopoda, 1282
soil water regimen, 1282
direct observations,
cotton, 1362
diet selection, 1362
habitat selection, 1362
omnivory, 1362
dispersal, 487, 644, 650, 1155, 1253, 1421, 1722
Anoplophora glabripennis, 644
Asian longhorned beetle, 644
Bt maize, 487
Cydia (Grapholita) molesta, 1155
density, 1421
Diabrotica virgifera virgifera, 650
diel periodicity, 650
elemental marking, 487
flight, 650
flight behavior, 1253
harmonic radar, 644
individual mark–recapture, 644
Leptinotarsa decemlineata, 1421
Loxostege sticticalis, 1253
mark-release-recapture, 1421
metapopulation biology, 1722
migration, 1253
oviposition preference, 1722
photoperiod, 1155
preimaginal environment, 1155
radar observation, 1253
Solanum linnaeanum, 1722
Solanum torvum, 1722
southwestern corn borer, 487
temperature, 1155
western corn rootworm, 650
dispersal-reproduction syndrome,
energy reserves, 1471
host plant quality, 1471
Phthorimaea operculella, 1471
wing loading, 1471
disruptants,
nonhost volatiles, 221
Tomicus piniperda, 221
verbenone, 221
distribution,
Cryptolestes ferrugineus, 1003
movement, 1003
stored grain, 1003
temperature gradient, 1003
wandering, 1003
dung beetle,
assemblage composition, 45
Scarabaeoidea, 45
spatial variability, 45
trophic resource, 45
dung beetles, 188, 370
cattle dung, 370
horse dung, 370
nesting, 188
olfactometer, 370
ovarian condition, 188
resource availability, 188
sexual maturation, 188
trophic preferences, 370

E

ectoparasitoid,
Calpodes ethlius, 227
Elachertus scutellatus, 227
koinobiont, 227
synovigenic, 227
ectoparasitoids,
biological control, 1313
fruit fly parasitoids, 1313
hyperparasitism, 1313
nontarget impact, 1313
edge effects,
Cyclosa, 1296
Hawaii, 1296
lava flows, 1296
Tetragnatha, 1296
egg hatch,
model, 107
pea, 107
pea leaf weevil, 107
temperature-dependent development, 107
egg load,
flight behavior, 1389
flight mill, 1389
tarnished plant bug, 1389
vertical flight chamber, 1389
western tarnished plant bug, 1389
egg parasitoids,
premating dispersion, 855
Trichogramma, 855
virgin females, 855
Elachertus scutellatus,
Calpodes ethlius, 227
ectoparasitoid, 227
koinobiont, 227
synovigenic, 227
elemental marking, 487, 930
Bt maize, 487
corn, 930
dispersal, 487
resistance management, 930
southwestern corn borer, 487, 930
Zea mays, 930
elemol,
catnip, 1562
nepetalactone, 1562
osage orange, 1562
repellency, 1562
emergence,
degree-day, 1541
flight activity, 1541
Lissorhoptrus oryzophilus, 1541
Oryza sativa, 1541

Empoasca fabae, 890, 1381
alfalfa, 890
Medicago sativa, 890
olfactory cues, 1381
preference, 1381
proportion maps, 890
reflectance, 1381
temporal spatial dynamics, 890
Encarsia citrina,
developmental rate, 1185
euonymus scale, 1185
life table, 1185
Quadraspidiotus perniciosus, 1185
energy reserves,
dispersal-reproduction syndrome, 1471
host plant quality, 1471
Phthorimaea operculella, 1471
wing loading, 1471
Enoclerus lecontei,
biological control, 1554
chemical ecology, 1554
Enoclerus sphegeus, 1554
Temnochila chlorodia, 1554
Enoclerus sphegeus,
biological control, 1554
chemical ecology, 1554
Enoclerus lecontei, 1554
Temnochila chlorodia, 1554
entomopathogen,
fungal pathogen, 975
microbial control, 975
Solenopsis invicta, 975
entomopathogenic algae,
Helicosporidium, 54
Heliocoverpa zea, 54
Spodoptera exigua, 54
Trichoplusia ni, 54
entomopathogenic fungi, 62, 1127
Asian longhorned beetle, 62
Beauveria bassiana, 62
Beauveria brongniartii, 62
biological control, 1127
Malacosoma disstria, 1127
Metarhizium anisopliae, 62
soil moisture, 1127
temperature, 1127
environmental stress,
Diptera, 1282
flooding, 1282
Isopoda, 1282
soil water regimen, 1282
epidemiology,
beech bark disease, 1274
beech scale, 1274
Cryptococcus fagisuga, 1274
Nectria, 1274
eradication,
Anoplophora glabripennis, 435
invasive species, 435
mark–recapture, 435
spatial autocorrelation, 435
Euhrychiopsis lecontei,
biocontrol, 872
development time, 872
watermilfoil, 872
euonymus scale,
developmental rate, 1185
Encarsia citrina, 1185
life table, 1185
Quadraspidiotus perniciosus, 1185
Euphorbia esula,
Aphthona lacertosa, 147
Aphthona nigriscutis, 147
biological control, 147
developmental rates, 147
leafy spurge, 147
European corn borer,
natural enemies, 320
Ostrinia nubilalis, 320
vegetative diversity, 320
weeds, 320
Euura sp.,
plant vigor, 700
population dynamics, 700
preference-performance linkage, 700
Salix scouleriana, 700
Evania appendigaster,
freezing, 1321
gamma irradiation, 1321
heating, 1321
Periplaneta americana, 1321
UV irradiation, 1321
external/internal marking,
behavior, 362
host quality, 362
superparasitism, 362

F

fall armyworm, 881, 982
Bacillus thuringiensis, 982
corn, 982
host strains, 881
nucleopolyhedrovirus, 982
population dynamics, 881
Spodoptera frugiperda, 881
transgenic plants, 982
fall generation,
cold hardiness, 113
Locusta migratoria, 113
postwinter population, 113
summer generation, 113
fecundity, 128, 138
biological control, 128
Cerambycidae, 138
life history parameters, 128
longevity, 128, 138
mating status, 128
Phoracantha recurva, 138
Phoracantha semipunctata, 138
feeding, 860, 990
behavior, 860
crop rotation resistance, 860
Diabrotica virgifera virgifera, 860
mile-a-minute weed, 990
oviposition, 860
Polygonum perfoliatum, 990
Rhinoncomimus latipes, 990
simulated herbivory, 990
feeding punctures,
Anthonomus grandis grandis, 443
boll weevil, 443
cotton fruit sizes, 443
oviposition punctures, 443
feeding state,
Carabidae, 1442
forest, 1442
male-to-female ratio, 1442
movement patterns, 1442
radar tracking, 1442
feltleaf willow,
Cytospora, 21
Scolytidae, 21
Trypophloeus, 21
feral colony,
Apis mellifera, 727
coastal prairie, 727
honey bee, 727
pollen, 727
fertile islands,
microbes, 599
nitrogen, 599
shrubs, 599
spatial heterogeneity, 599
field corn,
Gryllus spp., 577
Harpalus pensylvanicus, 577
Lycosidae, 577
weeds, 577
field sex ratio,
Ageniaspis fuscicollis, 528
The Netherlands, 528
Yponomeuta, 528
fire ant,
biological control, 340
pathogen, 340
polygyny, 340
Solenopsis invicta, 340
fire damage,
Dendroctonus ponderosae, 1070
habitat selection, 1070
population density, 1070
flight,
Diabrotica virgifera virgifera, 650
diel periodicity, 650
dispersal, 650
western corn rootworm, 650

flight activity, 1541, 1689
degree-day, 1541
emergence, 1541
grapevine, 1689
Lissorhoptrus oryzophilus, 1541
nearctic leafhopper, 1689
Oryza sativa, 1541
phytoplasma vector, 1689
flight behavior, 1253, 1389
dispersal, 1253
egg load, 1389
flight mill, 1389
Loxostege sticticalis, 1253
migration, 1253
radar observation, 1253
tarnished plant bug, 1389
vertical flight chamber, 1389
western tarnished plant bug, 1389
flight mill,
egg load, 1389
flight behavior, 1389
tarnished plant bug, 1389
vertical flight chamber, 1389
western tarnished plant bug, 1389
flooding,
Diptera, 1282
environmental stress, 1282
Isopoda, 1282
soil water regimen, 1282
floral visitors,
Asteraceae, 297
similarity indexes, 297
temporal variation, 297
xeric community, 297
Florida Everglades restoration,
biological control weeds, 997
predictive studies, 997
food conversion efficiency,
invasive species, 1169
nutritional ecology, 1169
phytosterols, 1169
root feeding, 1169
western corn rootworm, 1169
foraging, 1020, 1602
mud puddling, 1020
Pieris napi, 1020
polygyne, 1602
red imported fire ant, 1602
resource partitioning, 1602
sodium, 1020
Solenopsis invicta, 1602
forecasting,
geographic information system, 257
Markov chain, 257
pest management, 257
spatial processes, 257
Forelius,
biological control, 587
Monomorium, 587
Pheidole, 587
S. molesta, 587
forest,
Carabidae, 1442
feeding state, 1442
male-to-female ratio, 1442
movement patterns, 1442
radar tracking, 1442
forest fire,
boreal forest, 1703
habitat location, 1703
host selection, 1703
Monochamus scutellatus, 1703
Formosan subterranean termite,
monitoring station, 242
native subterranean termite, 242
termite baiting system, 242
termite foraging activity, 242
fragmentation,
habitat loss, 234
matrix-effects, 234
Ostrinia nubilalis, 234
Plathypena scabra, 234
species replacement, 234
freezing,
Evania appendigaster, 1321
gamma irradiation, 1321
heating, 1321
Periplaneta americana, 1321
UV irradiation, 1321
fruit fly parasitoids,
biological control, 1313
ectoparasitoids, 1313
hyperparasitism, 1313
nontarget impact, 1313
fungal pathogen,
entomopathogen, 975
microbial control, 975
Solenopsis invicta, 975
fungicides,
amorphogenic effects, 1151
Apis mellifera, 1151
honey bee, 1151
toxic effects, 1151

G

Galerucella calmariensis,
biological control, 356
Lythrum salicaria, 356
predation, 356
gall,
Asphondylia atriplicis, 709
Atriplex canescens, 709
Cecidomyiidae, 709
parasitoids, 709
gamma irradiation,
Evania appendigaster, 1321
freezing, 1321
heating, 1321
Periplaneta americana, 1321
UV irradiation, 1321
garlic mustard,
Carabidae, 564
geostatistics, 564
impact, 564
invasive, 564
gas exchange,
alfalfa, 450
herbivory, 450
Medicago spp., 450
red clover, 450
gene flow, 457, 1223
Anthonomus grandis, 457
genetic differentiation, 1223
genetic diversity, 457
Heliothis virescens, 1223
mitochondrial DNA restriction fragment length polymorphism, 457
phylogeography, 457
population genetics, 1223
RAPD-PCR markers, 1223
generalist,
adaptation, 1477
oviposition, 1477
polyphagy, 1477
specialist, 1477
generalist predator,
big-eyed bug, 964
Helicoverpa zea, 964
life table, 964
Spodoptera exigua, 964
generalist predators,
biological control, 662
predator-prey interactions, 662
prey vulnerability, 662
genetic differentiation,
gene flow, 1223
Heliothis virescens, 1223
population genetics, 1223
RAPD-PCR markers, 1223
genetic diversity,
Anthonomus grandis, 457
gene flow, 457
mitochondrial DNA restriction fragment length polymorphism, 457
phylogeography, 457
genotype,
apple maggot, 1695
host finding, 1695
ovipositional boring, 1695
preference, 1695
Geocoris,
aphids, 1014
Coleomegilla, 1014
Orius, 1014
predators, 1014
twospotted spider mite, 1014
geographic information system,
forecasting, 257
Markov chain, 257
pest management, 257

spatial processes, 257
geographic variation,
aphid, 776
biological control, 776
cold tolerance, 776
parasitoid, 776
geostatistics,
Carabidae, 564
garlic mustard, 564
impact, 564
invasive, 564
glassy-winged sharpshooter, 88, 165
amino acids, 165
dietary nitrogen, 165
diffusion model, 88
Homalodisca coagulata, 165
IgG protein markers, 88
insect rearing, 165
smoke tree sharpshooter, 88
yellow sticky traps, 88
global change,
CO_2 enrichment, 1609
insect communities, 1609
invasive species, 1609
plant communities, 1609
glucose,
Aphis gossypii, 690
cotton, 690
cotton aphid, 690
leaf sugars, 690
sucrose, 690
grain sorghum,
corn earworm, 1371
cotton, 1371
Helicoverpa zea, 1371
trap crop, 1371
grandisoic acid,
benzaldehyde, 389
Conotrachelus nenuphar, 389
monitoring traps, 389
plum curculio, 389
grapevine,
flight activity, 1689
nearctic leafhopper, 1689
phytoplasma vector, 1689
Grapholita,
lingonberry fruitworm, 405
pheromone traps, 405
population monitoring, 405
Vaccinium, 405
Grapholita molesta,
attract kill, 213
attracticide, 213
Oriental fruit moth, 213
pheromone, 213
Grapholita prunivora,
lesser apple worm, 506
Tortricidae, 506
grass flies,
anautogenous, 12
hexyl butanoate, 12
kairomone, 12
Olcella, 12
grass webworm,
developmental polymorphism, 1178
head capsule widths, 1178
grasshoppers,
hatch rate, 1528
overwintering egg, 1528
postdiapause development, 1528
grassy weeds,
alternate hosts, 1497
Diabrotica barberi, 1497
host-searching behavior, 1497
northern corn rootworm, 1497
green lacewings,
augmentative biological control, 1652
balsam fir, 1652
Christmas trees, 1652
Chrysoperla rufilabris, 1652
gregarization,
aggregation, 1577
phase, 1577
Schistocerca gregaria, 1577
Gryllus spp.,
field corn, 577
Harpalus pensylvanicus, 577
Lycosidae, 577
weeds, 577
gypsy moth,
Lepidoptera, 418
population dynamics, 418
sampling techniques, 418

H

habitat,
Ixodes scapularis, 900
landscape change, 900
habitat distribution,
Ixodes scapularis, 1266
spatial distribution, 1266
survival, 1266
habitat location,
boreal forest, 1703
forest fire, 1703
host selection, 1703
Monochamus scutellatus, 1703
habitat loss,
fragmentation, 234
matrix-effects, 234
Ostrinia nubilalis, 234
Plathypena scabra, 234
species replacement, 234
habitat manipulation,
conservation biological control, 1662
turf, 1662
habitat selection, 1070, 1362
cotton, 1362
Dendroctonus ponderosae, 1070
diet selection, 1362
direct observations, 1362
fire damage, 1070
omnivory, 1362
population density, 1070
habitat temperature,
Coptotermes formosanus, 197
Reticulitermes flavipes, 197
seasonal variation, 197
thermal tolerances, 197
hard-locked,
boll rot, 1095
cotton, 1095
stink bugs, 1095
Harmonia axyridis,
cold hardiness, 815
low temperature, 815
overwintering, 815
supercooling point, 815
harmonic radar,
Anoplophora glabripennis, 644
Asian longhorned beetle, 644
dispersal, 644
individual mark–recapture, 644
Harpalus pensylvanicus,
field corn, 577
Gryllus spp., 577
Lycosidae, 577
weeds, 577
hatch rate,
grasshoppers, 1528
overwintering egg, 1528
postdiapause development, 1528
Hawaii,
Cyclosa, 1296
edge effects, 1296
lava flows, 1296
Tetragnatha, 1296
head capsule widths,
developmental polymorphism, 1178
grass webworm, 1178
heating,
Evania appendigaster, 1321
freezing, 1321
gamma irradiation, 1321
Periplaneta americana, 1321
UV irradiation, 1321
Helicosporidium,
entomopathogenic algae, 54
Heliocoverpa zea, 54
Spodoptera exigua, 54
Trichoplusia ni, 54
Helicoverpa armigera,
host plants, 1570
life table, 1570
Helicoverpa zea, 964, 1371
big-eyed bug, 964
corn earworm, 1371
cotton, 1371
generalist predator, 964

grain sorghum, 1371
life table, 964
Spodoptera exigua, 964
trap crop, 1371
Heliocoverpa zea,
entomopathogenic algae, 54
Helicosporidium, 54
Spodoptera exigua, 54
Trichoplusia ni, 54
Heliothis virescens, 633, 1223
ascovirus transmission, 633
Campoletis sonorensis, 633
Cardiochiles nigriceps, 633
gene flow, 1223
genetic differentiation, 1223
Microplitis croceipes, 633
population genetics, 1223
RAPD-PCR markers, 1223
herbivorous mites,
induced susceptibility, 1088
papaya, 1088
powdery mildew, 1088
herbivory, 450, 1351
alfalfa, 450
Christmas berry, 1351
gas exchange, 450
leafroller, 1351
life history, 1351
Medicago spp., 450
red clover, 450
weed biological control, 1351
hexyl butanoate,
anautogenous, 12
grass flies, 12
kairomone, 12
Olcella, 12
Hippodamia convergens,
biological control, 282
conservation, 282
Scymnus loewii, 282
Homalodisca coagulata,
amino acids, 165
dietary nitrogen, 165
glassy-winged sharpshooter, 165
insect rearing, 165
honey bee, 378, 727, 1151
amorphogenic effects, 1151
Apis mellifera, 727, 1151
coastal prairie, 727
feral colony, 727
fungicides, 1151
insecticides, 378
learning, 378
Pavlovian conditioning, 378
pollen, 727
proboscis extension, 378
toxic effects, 1151
horse dung,
cattle dung, 370
dung beetles, 370
olfactometer, 370
trophic preferences, 370
host abundance,
Microctonus aethiopoides Loan, 941
modeling, 941
nontarget parasitism, 941
host aggregation,
parasitism, 1327
host density,
host mortality, 328
progeny production, 328
sex ratio, 328
host finding, 1410, 1695
Anthonomus pomorum, 1410
apple maggot, 1695
colors, 1410
genotype, 1695
host plant, 1410
ovipositional boring, 1695
preference, 1695
vision, 1410
host location,
chromatic contrast, 81
pupal parasitoid, 81
vision, 81
wavelength discrimination, 81
host modification,
Acyrthosiphon pisum, 1523
aphid, 1523
Aphidius ervi, 1523
host–parasitoid interactions, 1523
host mortality,
host density, 328
progeny production, 328
sex ratio, 328
host plant, 756, 1137, 1410
Anthonomus pomorum, 1410
canola, 756
colors, 1410
host finding, 1410
larval development, 756
light phase, 1137
Macrolophus pygmaeus, 1137
polyphagous, 756
predation rate, 1137
temperature, 1137
vision, 1410
host plant quality,
dispersal-reproduction syndrome, 1471
energy reserves, 1471
Phthorimaea operculella, 1471
wing loading, 1471
host plant resistance, 748, 1207
coat protein–mediated resistance, 1207
coevolution, 748
deme formation, 748
host specialization, 748
insect behavior, 1207
local adaptation, 748
virus vectors, 1207
virus-induced volatiles, 1207
host plants,
Helicoverpa armigera, 1570
life table, 1570
host quality, 362, 1488
behavior, 362
external/internal marking, 362
hyperparasitoid, 1488
idiobiont, 1488
parasitoid, 1488
superparasitism, 362
host range,
biocontrol, 1436
Solenopsis geminata, 1436
Solenopsis invicta, 1436
Solenopsis xyloni, 1436
host selection, 520, 1703
biological control, 520
boreal forest, 1703
forest fire, 1703
habitat location, 1703
Monochamus scutellatus, 1703
parasitoid, 520
Scarabaeidae, 520
turfgrass, 520
host specialization,
coevolution, 748
deme formation, 748
host plant resistance, 748
local adaptation, 748
host strains,
fall armyworm, 881
population dynamics, 881
Spodoptera frugiperda, 881
host volatiles,
attractants, 844
Cerambycidae, 844
Tetropium fuscum, 844
traps, 844
host–parasitoid interactions,
Acyrthosiphon pisum, 1523
aphid, 1523
Aphidius ervi, 1523
host modification, 1523
host-searching behavior,
alternate hosts, 1497
Diabrotica barberi, 1497
grassy weeds, 1497
northern corn rootworm, 1497
house fly,
biological control, 823
cold storage, 823
pteromalids, 823
survival, 823
hoverflies,
light-emitting diodes, 1416
thrips, 1416
traps, 1416
hybridization,
Aphis fabae, 1081
Cirsium arvense, 1081

clones, 1081
plant species preference, 1081
hyperparasitism,
biological control, 1313
ectoparasitoids, 1313
fruit fly parasitoids, 1313
nontarget impact, 1313
hyperparasitoid,
host quality, 1488
idiobiont, 1488
parasitoid, 1488
hypoxia,
caterpillar ecology, 1535
Coenonympha tullia nipisiquit, 1535
salt marsh, 1535
tidal submergence, 1535

I

idiobiont,
host quality, 1488
hyperparasitoid, 1488
parasitoid, 1488
IgG protein markers,
diffusion model, 88
glassy-winged sharpshooter, 88
smoke tree sharpshooter, 88
yellow sticky traps, 88
immigration,
CO potato beetle, 267
colonization, 267
Leptinotarsa decemlineata, 267
mapping, 267
impact,
Carabidae, 564
garlic mustard, 564
geostatistics, 564
invasive, 564
impatiens,
inoculative augmentation, 1344
nematode rearing, 1344
spinosad, 1344
western flower thrips, 1344
imported fire ants,
photointerpretation, 1718
remote sensing, 1718
inbreeding depression,
biological control, 765
Nasonia vitripennis, 765
outbreeding depression, 765
parasitoid, 765
individual mark–recapture,
Anoplophora glabripennis, 644
Asian longhorned beetle, 644
dispersal, 644
harmonic radar, 644
induced susceptibility,
herbivorous mites, 1088
papaya, 1088
powdery mildew, 1088
inoculative augmentation,
impatiens, 1344
nematode rearing, 1344
spinosad, 1344
western flower thrips, 1344
insect behavior, 1025, 1207
Bacillus thuringiensis, 1025
coat protein–mediated resistance, 1207
diamondback moth, 1025
host plant resistance, 1207
Plutella xylostella, 1025
transgenic crop, 1025
virus vectors, 1207
virus-induced volatiles, 1207
insect communities,
CO_2 enrichment, 1609
global change, 1609
invasive species, 1609
plant communities, 1609
insect ecology,
detritivore, 907
ordination, 907
webspinners, 907
insect rearing,
amino acids, 165
dietary nitrogen, 165
glassy-winged sharpshooter, 165
Homalodisca coagulata, 165
insect vector,
morphological variation, 1192
Zea mays, 1192
insect–fungal interaction,
bark beetles, 28
Ophiostoma ips, 28
ophiostomatoid fungi, 28
symbiosis, 28
insecticides,
honey bee, 378
learning, 378
Pavlovian conditioning, 378
proboscis extension, 378
insects,
potato, 1632
straw mulch, 1632
watermelon, 1632
weeds, 1632
insect–weed interactions,
Lissorhoptrus oryzophilus, 720
Oebalus pugnax, 720
Oryza sativa, 720
interactions,
Lacanobia oleracea, 949
Lacanobia oleracea granulovirus, 949
Meteorus gyrator, 949
introduced predator,
antipredator behavior, 1032
biological control, 1032
Pardosa milvina, 1032
Tenodera aridifolia sinensis, 1032
introduced species,
deme extinction, 182
mantids, 182
metapopulations, 182
Tenodera aridifolia sinensis, 182
invasive,
Carabidae, 564
garlic mustard, 564
geostatistics, 564
impact, 564
invasive species, 206, 435, 540, 608, 1169, 1609
Acer mono, 540
Anoplophora glabripennis, 435, 540
Asian longhorned beetle, 540
biological control, 608
CO_2 enrichment, 1609
eradication, 435
food conversion efficiency, 1169
global change, 1609
insect communities, 1609
mark–recapture, 435
monoterpenes, 206
nutritional ecology, 1169
phylogenetic geography, 206
phytosterols, 1169
plant communities, 1609
predator community, 608
root feeding, 1169
semiochemicals, 206
South Korea, 540
spatial autocorrelation, 435
western corn rootworm, 1169
Isopoda,
Diptera, 1282
environmental stress, 1282
flooding, 1282
soil water regimen, 1282
isothermal calorimetry,
multicolored Asian lady beetle, 832
R_{CO2}, 832
respiratory physiology, 832
temperature dependence, 832
Ixodes scapularis, 900, 1266
habitat, 900
habitat distribution, 1266
landscape change, 900
spatial distribution, 1266
survival, 1266

K

kairomone,
anautogenous, 12
grass flies, 12
hexyl butanoate, 12
Olcella, 12

koinobiont,
Calpodes ethlius, 227
ectoparasitoid, 227
Elachertus scutellatus, 227
synovigenic, 227
Konza prairie,
bison grazing, 1617
Orthoptera, 1617
population dynamics, 1617
prescribed burning, 1617

L

Lacanobia oleracea,
interactions, 949
Lacanobia oleracea granulovirus, 949
Meteorus gyrator, 949
Lacanobia oleracea granulovirus,
interactions, 949
Lacanobia oleracea, 949
Meteorus gyrator, 949
landscape change,
habitat, 900
Ixodes scapularis, 900
landscape ecology,
Acyrthosiphon pisum, 1593
Aphis glycines, 1593
Rhopalosiphum maidis, 1593
Therioaphis trifolii, 1593
larval development,
canola, 756
host plant, 756
polyphagous, 756
latitudinal variation,
acclimation, 155
cold hardiness, 155
Liriomyza huidobrensis, 155
pea leafminer, 155
lava flows,
Cyclosa, 1296
edge effects, 1296
Hawaii, 1296
Tetragnatha, 1296
leaf sugars,
Aphis gossypii, 690
cotton, 690
cotton aphid, 690
glucose, 690
sucrose, 690
leafroller,
Christmas berry, 1351
herbivory, 1351
life history, 1351
weed biological control, 1351
leafy spurge,
Aphthona lacertosa, 147
Aphthona nigriscutis, 147
biological control, 147
developmental rates, 147
Euphorbia esula, 147
learning,
honey bee, 378
insecticides, 378
Pavlovian conditioning, 378
proboscis extension, 378
Lepidoptera,
gypsy moth, 418
population dynamics, 418
sampling techniques, 418
Leptinotarsa decemlineata, 267, 275, 1421
biotechnology, 275
CO potato beetle, 267
colonization, 267
Cry3Aa, 275
density, 1421
dispersal, 1421
immigration, 267
mapping, 267
mark-release-recapture, 1421
nontarget effects, 275
potato pest management, 275
lesser apple worm,
Grapholita prunivora, 506
Tortricidae, 506
life history, 807, 1200, 1351
Christmas berry, 1351
corn, 807
herbivory, 1351
leafroller, 1351
life table, 1200
modeling, 807
Podisus maculiventris, 1200
Sitotroga cerealella, 807
survivorship, 807
weed biological control, 1351
life history parameters,
biological control, 128
fecundity, 128
longevity, 128
mating status, 128
life table, 964, 1185, 1200, 1570
big-eyed bug, 964
developmental rate, 1185
Encarsia citrina, 1185
euonymus scale, 1185
generalist predator, 964
Helicoverpa armigera, 1570
Helicoverpa zea, 964
host plants, 1570
life history, 1200
Podisus maculiventris, 1200
Quadraspidiotus perniciosus, 1185
Spodoptera exigua, 964
life table statistics,
Phthorimaea operculella, 477
population dynamics, 477
potato pests, 477
potato tuber moth, 477
light phase,
host plant, 1137
Macrolophus pygmaeus, 1137
predation rate, 1137
temperature, 1137
light-emitting diodes,
hoverflies, 1416
thrips, 1416
traps, 1416
lingonberry fruitworm,
Grapholita, 405
pheromone traps, 405
population monitoring, 405
Vaccinium, 405
Liriomyza huidobrensis,
acclimation, 155
cold hardiness, 155
latitudinal variation, 155
pea leafminer, 155
Lissorhoptrus oryzophilus, 720, 1541
degree-day, 1541
emergence, 1541
flight activity, 1541
insect–weed interactions, 720
Oebalus pugnax, 720
Oryza sativa, 720, 1541
local adaptation,
coevolution, 748
deme formation, 748
host plant resistance, 748
host specialization, 748
Locusta migratoria,
cold hardiness, 113
fall generation, 113
postwinter population, 113
summer generation, 113
longevity, 128, 138, 779
biological control, 128
Cerambycidae, 138
costs of mating, 779
fecundity, 128, 138
life history parameters, 128
mating status, 128
Phoracantha recurva, 138
Phoracantha semipunctata, 138
repeatability, 779
sexual selection, 779
starvation resistance, 779
low temperature,
cold hardiness, 815
Harmonia axyridis, 815
overwintering, 815
supercooling point, 815
Loxostege sticticalis,
dispersal, 1253
flight behavior, 1253
migration, 1253
radar observation, 1253
Lycosidae,
field corn, 577
Gryllus spp., 577

Harpalus pensylvanicus, 577
weeds, 577
Lygus elisus,
developmental duration, 1549
population dynamics, 1549
temperature, 1549
Lythrum salicaria,
biological control, 356
Galerucella calmariensis, 356
predation, 356

M

Macrolophus pygmaeus, 499, 1137
biological control, 499
host plant, 1137
light phase, 1137
Miridae, 499
Myzus persicae, 499
pepper, 499
predation rate, 1137
temperature, 1137
Malacosoma disstria,
biological control, 1127
entomopathogenic fungi, 1127
soil moisture, 1127
temperature, 1127
male aggression,
body size, 657
mating success, 657
Nadezhdiella cantori, 657
sexual selection, 657
male-to-female ratio,
Carabidae, 1442
feeding state, 1442
forest, 1442
movement patterns, 1442
radar tracking, 1442
mantids,
deme extinction, 182
introduced species, 182
metapopulations, 182
Tenodera aridifolia sinensis, 182
mapping,
CO potato beetle, 267
colonization, 267
immigration, 267
Leptinotarsa decemlineata, 267
Markov chain,
forecasting, 257
geographic information system, 257
pest management, 257
spatial processes, 257
mark–recapture,
Anoplophora glabripennis, 435
eradication, 435
invasive species, 435
spatial autocorrelation, 435
mark-release-recapture,
density, 1421
dispersal, 1421
Leptinotarsa decemlineata, 1421
mating status,
biological control, 128
fecundity, 128
life history parameters, 128
longevity, 128
mating success,
body size, 657
male aggression, 657
Nadezhdiella cantori, 657
sexual selection, 657
matrix-effects,
fragmentation, 234
habitat loss, 234
Ostrinia nubilalis, 234
Plathypena scabra, 234
species replacement, 234
Medicago sativa, 119, 890
alfalfa, 119, 890
clover root curculio, 119
Empoasca fabae, 890
nodules, 119
proportion maps, 890
Sitona hispidulus, 119
temporal spatial dynamics, 890
Medicago spp.,
alfalfa, 450
gas exchange, 450
herbivory, 450
red clover, 450
metapopulation biology,
dispersal, 1722
oviposition preference, 1722
Solanum linnaeanum, 1722
Solanum torvum, 1722
metapopulations,
deme extinction, 182
introduced species, 182
mantids, 182
Tenodera aridifolia sinensis, 182
Metarhizium anisopliae, 62, 1335
Asian longhorned beetle, 62
Bacillus thuringiensis, 1335
Beauveria bassiana, 62, 1335
Beauveria brongniartii, 62
entomopathogenic fungi, 62
soil survey, 1335
Steinernema oregonense, 1335
Meteorus gyrator,
interactions, 949
Lacanobia oleracea, 949
Lacanobia oleracea granulovirus, 949
microbes,
fertile islands, 599
nitrogen, 599
shrubs, 599
spatial heterogeneity, 599
microbial control,
entomopathogen, 975
fungal pathogen, 975
Solenopsis invicta, 975
Microctonus aethiopoides Loan,
host abundance, 941
modeling, 941
nontarget parasitism, 941
Microplitis croceipes,
ascovirus transmission, 633
Campoletis sonorensis, 633
Cardiochiles nigriceps, 633
Heliothis virescens, 633
migration, 249, 1253
dispersal, 1253
flight behavior, 1253
Loxostege sticticalis, 1253
monarchs, 249
radar observation, 1253
stopover, 249
timing, 249
weather, 249
mile-a-minute weed,
feeding, 990
Polygonum perfoliatum, 990
Rhinoncomimus latipes, 990
simulated herbivory, 990
Miridae,
biological control, 499
Macrolophus pygmaeus, 499
Myzus persicae, 499
pepper, 499
mitochondrial DNA restriction fragment length polymorphism,
Anthonomus grandis, 457
gene flow, 457
genetic diversity, 457
phylogeography, 457
model, 1, 107, 787
development, 1
egg hatch, 107
Nephus bisignatus, *1*
Nephus includens, 1
Ostrinia furnacalis, 787
oviposition behavior, 787
pea, 107
pea leaf weevil, 107
temperature-dependent development, 107, 787
threshold, 1
Trichogramma ostriniae, 787
modeling, 426, 807, 941
corn, 807
decision support tools, 426
host abundance, 941
life history, 807
Microctonus aethiopoides Loan, 941
nontarget parasitism, 941
Oklahoma, 426
Sitotroga cerealella, 807

stored-product beetles, 426
survivorship, 807
mole cricket,
acoustic communication, 1431
moon phase, 1431
sex ratio, 1431
Monarch butterflies,
Asclepias syriaca, 1062
compensatory feeding, 1062
Danaus plexippus, 1062
plant nitrogen, 1062
monarch butterfly,
Bt corn, 1116
chronic effects, 1116
nontarget, 1116
risk assessment, 1116
monarchs,
migration, 249
stopover, 249
timing, 249
weather, 249
monitoring,
Oklahoma, 1465
stored-product beetles, 1465
trap catch analysis, 1465
monitoring station,
Formosan subterranean termite, 242
native subterranean termite, 242
termite baiting system, 242
termite foraging activity, 242
monitoring traps,
benzaldehyde, 389
Conotrachelus nenuphar, 389
grandisoic acid, 389
plum curculio, 389
Monochamus scutellatus,
boreal forest, 1703
forest fire, 1703
habitat location, 1703
host selection, 1703
Monomorium,
biological control, 587
Forelius, 587
Pheidole, 587
S. molesta, 587
monoterpenes,
invasive species, 206
phylogenetic geography, 206
semiochemicals, 206
moon phase,
acoustic communication, 1431
mole cricket, 1431
sex ratio, 1431
morphological variation,
insect vector, 1192
Zea mays, 1192
mounds,
multispectral, 1045
remote sensing, 1045
movement,
Cryptolestes ferrugineus, 1003
distribution, 1003
stored grain, 1003
temperature gradient, 1003
wandering, 1003
movement patterns,
Carabidae, 1442
feeding state, 1442
forest, 1442
male-to-female ratio, 1442
radar tracking, 1442
mud puddling,
foraging, 1020
Pieris napi, 1020
sodium, 1020
multicolored Asian lady beetle,
isothermal calorimetry, 832
R_{CO2}, 832
respiratory physiology, 832
temperature dependence, 832
multispectral,
mounds, 1045
remote sensing, 1045
Myzus persicae,
biological control, 499
Macrolophus pygmaeus, 499
Miridae, 499
pepper, 499

N

Nadezhdiella cantori,
body size, 657
male aggression, 657
mating success, 657
sexual selection, 657
Nasonia vitripennis,
biological control, 765
inbreeding depression, 765
outbreeding depression, 765
parasitoid, 765
native ant,
predation, 1626
red imported fire ant, 1626
Solenopsis molesta, 1626
thief ant, 1626
native subterranean termite,
Formosan subterranean termite, 242
monitoring station, 242
termite baiting system, 242
termite foraging activity, 242
natural enemies,
European corn borer, 320
Ostrinia nubilalis, 320
vegetative diversity, 320
weeds, 320
natural habitats,
risk, 310
vineyards, 310
wild grape, 310
nearctic leafhopper,
flight activity, 1689
grapevine, 1689
phytoplasma vector, 1689
nectar robbing,
Apis mellifera, 100
rabbiteye blueberry, 100
Vaccinium ashei, 100
Xylocopa virginica, 100
Nectria,
beech bark disease, 1274
beech scale, 1274
Cryptococcus fagisuga, 1274
epidemiology, 1274
nematode rearing,
impatiens, 1344
inoculative augmentation, 1344
spinosad, 1344
western flower thrips, 1344
nepetalactone,
catnip, 1562
elemol, 1562
osage orange, 1562
repellency, 1562
Nephus bisignatus,
development, 1
model, 1
Nephus includens, 1
threshold, 1
Nephus includens,
development, 1
model, 1
Nephus bisignatus, 1
threshold, 1
nesting,
dung beetles, 188
ovarian condition, 188
resource availability, 188
sexual maturation, 188
nitrogen,
fertile islands, 599
microbes, 599
shrubs, 599
spatial heterogeneity, 599
nodules,
alfalfa, 119
clover root curculio, 119
Medicago sativa, 119
Sitona hispidulus, 119
nonhost volatiles,
disruptants, 221
Tomicus piniperda, 221
verbenone, 221
nontarget, 554, 921, 1116
Bacillus thuringiensis, 921
Bt corn, 1116
Bt cotton, 921
canonical correspondence analysis, 554
chronic effects, 1116
crop rotation, 554

Cry1Ab, 554
monarch butterfly, 1116
pink bollworm, 921
predators, 554
risk assessment, 1116
nontarget effects,
biotechnology, 275
Cry3Aa, 275
Leptinotarsa decemlineata, 275
potato pest management, 275
nontarget impact,
biological control, 1313
ectoparasitoids, 1313
fruit fly parasitoids, 1313
hyperparasitism, 1313
nontarget parasitism,
host abundance, 941
Microctonus aethiopoides Loan, 941
modeling, 941
nontargets,
Danaus plexippus, 1109
risk assessment, 1109
transgenic corn, 1109
northern corn rootworm,
alternate hosts, 1497
Diabrotica barberi, 1497
grassy weeds, 1497
host-searching behavior, 1497
nucleopolyhedrovirus,
Bacillus thuringiensis, 982
corn, 982
fall armyworm, 982
transgenic plants, 982
nutritional ecology,
food conversion efficiency, 1169
invasive species, 1169
phytosterols, 1169
root feeding, 1169
western corn rootworm, 1169

O

Oebalus pugnax,
insect–weed interactions, 720
Lissorhoptrus oryzophilus, 720
Oryza sativa, 720
Oklahoma, 426, 1465
decision support tools, 426
modeling, 426
monitoring, 1465
stored-product beetles, 426, 1465
trap catch analysis, 1465
Olcella,
anautogenous, 12
grass flies, 12
hexyl butanoate, 12
kairomone, 12
olfaction,
Oryzaephilus surinamensis, 75
packaging, 75
sawtoothed grain beetle, 75
stored products, 75
olfactometer,
cattle dung, 370
dung beetles, 370
horse dung, 370
trophic preferences, 370
olfactory cues,
Empoasca fabae, 1381
preference, 1381
reflectance, 1381
omnivory,
cotton, 1362
diet selection, 1362
direct observations, 1362
habitat selection, 1362
Ophiostoma ips,
bark beetles, 28
insect–fungal interaction, 28
ophiostomatoid fungi, 28
symbiosis, 28
ophiostomatoid fungi,
bark beetles, 28
insect–fungal interaction, 28
Ophiostoma ips, 28
symbiosis, 28
ordination,
detritivore, 907
insect ecology, 907
webspinners, 907
Oriental fruit moth,
attract kill, 213
attracticide, 213
Grapholita molesta, 213
pheromone, 213
Orius,
aphids, 1014
Coleomegilla, 1014
Geocoris, 1014
predators, 1014
twospotted spider mite, 1014
Orthoptera,
bison grazing, 1617
Konza prairie, 1617
population dynamics, 1617
prescribed burning, 1617
Oryza sativa, 720, 1541
degree-day, 1541
emergence, 1541
flight activity, 1541
insect–weed interactions, 720
Lissorhoptrus oryzophilus, 720, 1541
Oebalus pugnax, 720
Oryzaephilus surinamensis,
olfaction, 75
packaging, 75
sawtoothed grain beetle, 75
stored products, 75
osage orange,
catnip, 1562
elemol, 1562
nepetalactone, 1562
repellency, 1562
Ostrinia furnacalis,
model, 787
oviposition behavior, 787
temperature-dependent development, 787
Trichogramma ostriniae, 787
Ostrinia nubilalis, 234, 320
European corn borer, 320
fragmentation, 234
habitat loss, 234
matrix-effects, 234
natural enemies, 320
Plathypena scabra, 234
species replacement, 234
vegetative diversity, 320
weeds, 320
outbreeding depression,
biological control, 765
inbreeding depression, 765
Nasonia vitripennis, 765
parasitoid, 765
ovarian condition,
dung beetles, 188
nesting, 188
resource availability, 188
sexual maturation, 188
overwintering, 513, 815
Adoxophyes orana, 513
cold hardiness, 815
diapause termination, 513
Harmonia axyridis, 815
low temperature, 815
photoperiod, 513
supercooling point, 815
Tortricidae, 513
overwintering egg,
grasshoppers, 1528
hatch rate, 1528
postdiapause development, 1528
overwintering stage,
bruchid, 1163
diapause, 1163
seasonal adaptation, 1163
ovigeny,
predator fitness, 1401
prey preference, 1401
oviposition, 860, 1217, 1477
adaptation, 1477
artificial substrate, 1217
behavior, 860
crop rotation resistance, 860
Diabrotica virgifera virgifera, 860
feeding, 860
generalist, 1477
plant extracts, 1217
polyphagy, 1477
specialist, 1477

oviposition behavior,
model, 787
Ostrinia furnacalis, 787
temperature-dependent development, 787
Trichogramma ostriniae, 787
oviposition preference,
dispersal, 1722
metapopulation biology, 1722
Solanum linnaeanum, 1722
Solanum torvum, 1722
oviposition punctures,
Anthonomus grandis grandis, 443
boll weevil, 443
cotton fruit sizes, 443
feeding punctures, 443
ovipositional boring,
apple maggot, 1695
genotype, 1695
host finding, 1695
preference, 1695

P

packaging,
olfaction, 75
Oryzaephilus surinamensis, 75
sawtoothed grain beetle, 75
stored products, 75
papaya,
herbivorous mites, 1088
induced susceptibility, 1088
powdery mildew, 1088
papaya ring spot virus,
aphids, 1242
Bemisia tabaci, 1242
squash leaf curl virus, 1242
watermelon, 1242
parasitism,
host aggregation, 1327
parasitoid, 291, 520, 765, 776, 1488
aphid, 776
Asteraceae, 291
biological control, 520, 765, 776
cold tolerance, 776
geographic variation, 776
host quality, 1488
host selection, 520
hyperparasitoid, 1488
idiobiont, 1488
inbreeding depression, 765
Nasonia vitripennis, 765
outbreeding depression, 765
prairie, 291
predation, 291
Scarabaeidae, 520
Silphium, 291
stable isotope, 291
turfgrass, 520
parasitoids,
Asphondylia atriplicis, 709
Atriplex canescens, 709
Cecidomyiidae, 709
gall, 709
Pardosa milvina,
antipredator behavior, 1032
biological control, 1032
introduced predator, 1032
Tenodera aridifolia sinensis, 1032
pathogen,
biological control, 340
fire ant, 340
polygyny, 340
Solenopsis invicta, 340
Pavlovian conditioning,
honey bee, 378
insecticides, 378
learning, 378
proboscis extension, 378
pea,
egg hatch, 107
model, 107
pea leaf weevil, 107
temperature-dependent development, 107
pea leaf weevil,
egg hatch, 107
model, 107
pea, 107
temperature-dependent development, 107
pea leafminer,
acclimation, 155
cold hardiness, 155
latitudinal variation, 155
Liriomyza huidobrensis, 155
Pea protein,
Anisopteromalus calandrae, 671
Cephalonomia waterstoni, 671
Cryptolestes, 671
Sitophilus, 671
peach,
Conotrachelus nenuphar, 174
degree-day model, 174
plum curculio, 174
Prunus persica, 174
pepper,
biological control, 499
Macrolophus pygmaeus, 499
Miridae, 499
Myzus persicae, 499
Periplaneta americana,
Evania appendigaster, 1321
freezing, 1321
gamma irradiation, 1321
heating, 1321
UV irradiation, 1321
pest management,
forecasting, 257
geographic information system, 257
Markov chain, 257
spatial processes, 257
pest monitoring,
Argyrotaenia velutinana, 1711
codling moth, 1711
Cydia pomonella, 1711
redbanded leafroller, 1711
phase,
aggregation, 1577
gregarization, 1577
Schistocerca gregaria, 1577
Pheidole,
biological control, 587
Forelius, 587
Monomorium, 587
S. molesta, 587
phenological model,
subarctic, 1513
thermoregulation, 1513
phenology,
corn, 35
resistance, 35
rotation, 35
soybean, 35
pheromone,
attract kill, 213
attracticide, 213
Grapholita molesta, 213
Oriental fruit moth, 213
pheromone trapping,
Aesculus hippocastanum, 1584
biological invasion, 1584
Cameraria ohridella, 1584
spatial dynamics, 1584
pheromone traps, 397, 405
damage prediction, 397
Grapholita, 405
lingonberry fruitworm, 405
Pinus taeda, 397
population monitoring, 405
Rhyacionia frustrana, 397
Vaccinium, 405
Phoracantha recurva, 138, 1232
Cerambycidae, 138
competition, 1232
diapause, 1232
fecundity, 138
longevity, 138
Phoracantha semipunctata, 138, 1232
seasonal development, 1232
Phoracantha semipunctata, 138, 1232
Cerambycidae, 138
competition, 1232
diapause, 1232
fecundity, 138
longevity, 138

Phoracantha recurva, 138, 1232
seasonal development, 1232
photointerpretation,
imported fire ants, 1718
remote sensing, 1718
photoperiod, 513, 1155
Adoxophyes orana, 513
Cydia (Grapholita) molesta, 1155
diapause termination, 513
dispersal, 1155
overwintering, 513
preimaginal environment, 1155
temperature, 1155
Tortricidae, 513
Phthorimaea operculella, 477, 1471
dispersal-reproduction syndrome, 1471
energy reserves, 1471
host plant quality, 1471
life table statistics, 477
population dynamics, 477
potato pests, 477
potato tuber moth, 477
wing loading, 1471
phylogenetic geography,
invasive species, 206
monoterpenes, 206
semiochemicals, 206
phylogeography,
Anthonomus grandis, 457
gene flow, 457
genetic diversity, 457
mitochondrial DNA restriction fragment length polymorphism, 457
physiological saline solution,
anaerobic protists, 1145
protist counting techniques, 1145
Reticulitermes flavipes, 1145
symbiotic protists, 1145
phytoplasma vector,
flight activity, 1689
grapevine, 1689
nearctic leafhopper, 1689
phytosterols,
food conversion efficiency, 1169
invasive species, 1169
nutritional ecology, 1169
root feeding, 1169
western corn rootworm, 1169
Pieris napi,
foraging, 1020
mud puddling, 1020
sodium, 1020
pink bollworm,
Bacillus thuringiensis, 921
Bt cotton, 921
nontarget, 921
Pinus taeda,
damage prediction, 397
pheromone traps, 397
Rhyacionia frustrana, 397
plant communities,
CO_2 enrichment, 1609
global change, 1609
insect communities, 1609
invasive species, 1609
plant extracts,
artificial substrate, 1217
oviposition, 1217
plant nitrogen,
Asclepias syriaca, 1062
compensatory feeding, 1062
Danaus plexippus, 1062
Monarch butterflies, 1062
plant quality,
aquatic weed, 346
damage, 346
Pontederiaceae, 346
plant species preference,
Aphis fabae, 1081
Cirsium arvense, 1081
clones, 1081
hybridization, 1081
plant vigor,
Euura sp., 700
population dynamics, 700
preference-performance linkage, 700
Salix scouleriana, 700
plant-stress hypothesis,
Diplolepis spp., 1052
plant-vigor hypothesis, 1052
Rosa arkansana, 1052
tall-grass prairie, 1052
plant-vigor hypothesis,
Diplolepis spp., 1052
plant-stress hypothesis, 1052
Rosa arkansana, 1052
tall-grass prairie, 1052
Plathypena scabra,
fragmentation, 234
habitat loss, 234
matrix-effects, 234
Ostrinia nubilalis, 234
species replacement, 234
plum curculio, 174, 389
benzaldehyde, 389
Conotrachelus nenuphar, 174, 389
degree-day model, 174
grandisoic acid, 389
monitoring traps, 389
peach, 174
Prunus persica, 174
Plutella xylostella,
Bacillus thuringiensis, 1025
diamondback moth, 1025
insect behavior, 1025
transgenic crop, 1025
pod position,
canopy temperature, 471
defoliation, 471
Vigna unguiculata, 471
Podisus maculiventris,
life history, 1200
life table, 1200
pollen,
Apis mellifera, 727
coastal prairie, 727
feral colony, 727
honey bee, 727
pollinivory,
biological control, 958
risk assessment, 958
transgenic crops, 958
Zea mays, 958
Polygonum perfoliatum,
feeding, 990
mile-a-minute weed, 990
Rhinoncomimus latipes, 990
simulated herbivory, 990
polygyne,
foraging, 1602
red imported fire ant, 1602
resource partitioning, 1602
Solenopsis invicta, 1602
polygyny,
biological control, 340
fire ant, 340
pathogen, 340
Solenopsis invicta, 340
polyphagous,
canola, 756
host plant, 756
larval development, 756
polyphagy,
adaptation, 1477
generalist, 1477
oviposition, 1477
specialist, 1477
Pontederiaceae,
aquatic weed, 346
damage, 346
plant quality, 346
population density,
Dendroctonus ponderosae, 1070
fire damage, 1070
habitat selection, 1070
population dynamics, 418, 477, 700, 881, 1549, 1617
bison grazing, 1617
developmental duration, 1549
Euura sp., 700
fall armyworm, 881
gypsy moth, 418
host strains, 881
Konza prairie, 1617
Lepidoptera, 418
life table statistics, 477
Lygus elisus, 1549
Orthoptera, 1617

Phthorimaea operculella, 477
plant vigor, 700
potato pests, 477
potato tuber moth, 477
preference-performance linkage, 700
prescribed burning, 1617
Salix scouleriana, 700
sampling techniques, 418
Spodoptera frugiperda, 881
temperature, 1549
population genetics,
gene flow, 1223
genetic differentiation, 1223
Heliothis virescens, 1223
RAPD-PCR markers, 1223
population monitoring,
Grapholita, 405
lingonberry fruitworm, 405
pheromone traps, 405
Vaccinium, 405
postdiapause development,
grasshoppers, 1528
hatch rate, 1528
overwintering egg, 1528
postwinter population,
cold hardiness, 113
fall generation, 113
Locusta migratoria, 113
summer generation, 113
potato,
insects, 1632
straw mulch, 1632
watermelon, 1632
weeds, 1632
potato pest management,
biotechnology, 275
Cry3Aa, 275
Leptinotarsa decemlineata, 275
nontarget effects, 275
potato pests,
life table statistics, 477
Phthorimaea operculella, 477
population dynamics, 477
potato tuber moth, 477
potato tuber moth,
life table statistics, 477
Phthorimaea operculella, 477
population dynamics, 477
potato pests, 477
powdery mildew,
herbivorous mites, 1088
induced susceptibility, 1088
papaya, 1088
prairie,
Asteraceae, 291
parasitoid, 291
predation, 291
Silphium, 291
stable isotope, 291
prairie grasses,
alternate hosts, 740
Diabrotica virgifera virgifera, 740
resistance management, 740
predation, 291, 356, 1626
Asteraceae, 291
biological control, 356
Galerucella calmariensis, 356
Lythrum salicaria, 356
native ant, 1626
parasitoid, 291
prairie, 291
red imported fire ant, 1626
Silphium, 291
Solenopsis molesta, 1626
stable isotope, 291
thief ant, 1626
predation rate,
host plant, 1137
light phase, 1137
Macrolophus pygmaeus, 1137
temperature, 1137
predator community,
biological control, 608
invasive species, 608
predator fitness,
ovigeny, 1401
prey preference, 1401
predator-prey interactions,
biological control, 662
generalist predators, 662
prey vulnerability, 662
predators, 554, 1014
aphids, 1014
canonical correspondence analysis, 554
Coleomegilla, 1014
crop rotation, 554
Cry1Ab, 554
Geocoris, 1014
nontarget, 554
Orius, 1014
twospotted spider mite, 1014
predictive studies,
biological control weeds, 997
Florida Everglades restoration, 997
preference, 1381, 1695
apple maggot, 1695
Empoasca fabae, 1381
genotype, 1695
host finding, 1695
olfactory cues, 1381
ovipositional boring, 1695
reflectance, 1381
preference-performance linkage,
Euura sp., 700
plant vigor, 700
population dynamics, 700
Salix scouleriana, 700
preimaginal environment,
Cydia (Grapholita) molesta, 1155
dispersal, 1155
photoperiod, 1155
temperature, 1155
premating dispersion,
egg parasitoids, 855
Trichogramma, 855
virgin females, 855
prescribed burning,
bison grazing, 1617
Konza prairie, 1617
Orthoptera, 1617
population dynamics, 1617
prey preference,
ovigeny, 1401
predator fitness, 1401
prey vulnerability,
biological control, 662
generalist predators, 662
predator-prey interactions, 662
proboscis extension,
honey bee, 378
insecticides, 378
learning, 378
Pavlovian conditioning, 378
progeny production,
host density, 328
host mortality, 328
sex ratio, 328
proportion maps,
alfalfa, 890
Empoasca fabae, 890
Medicago sativa, 890
temporal spatial dynamics, 890
protist counting techniques,
anaerobic protists, 1145
physiological saline solution, 1145
Reticulitermes flavipes, 1145
symbiotic protists, 1145
Prunus persica,
Conotrachelus nenuphar, 174
degree-day model, 174
peach, 174
plum curculio, 174
pteromalids,
biological control, 823
cold storage, 823
house fly, 823
survival, 823
pupal parasitoid,
chromatic contrast, 81
host location, 81
vision, 81
wavelength discrimination, 81
pupation success,
Aethina tumida, 794
Apis mellifera, 794
soil properties, 794

Q

Quadraspidiotus perniciosus,
developmental rate, 1185
Encarsia citrina, 1185
euonymus scale, 1185
life table, 1185

R

R_{CO2},
isothermal calorimetry, 832
multicolored Asian lady beetle, 832
respiratory physiology, 832
temperature dependence, 832
rabbiteye blueberry,
Apis mellifera, 100
nectar robbing, 100
Vaccinium ashei, 100
Xylocopa virginica, 100
radar observation,
dispersal, 1253
flight behavior, 1253
Loxostege sticticalis, 1253
migration, 1253
radar tracking,
Carabidae, 1442
feeding state, 1442
forest, 1442
male-to-female ratio, 1442
movement patterns, 1442
RAPD-PCR markers,
gene flow, 1223
genetic differentiation, 1223
Heliothis virescens, 1223
population genetics, 1223
red clover,
alfalfa, 450
gas exchange, 450
herbivory, 450
Medicago spp., 450
red imported fire ant, 1602, 1626
foraging, 1602
native ant, 1626
polygyne, 1602
predation, 1626
resource partitioning, 1602
Solenopsis invicta, 1602
Solenopsis molesta, 1626
thief ant, 1626
redbanded leafroller,
Argyrotaenia velutinana, 1711
codling moth, 1711
Cydia pomonella, 1711
pest monitoring, 1711
reflectance,
Empoasca fabae, 1381
olfactory cues, 1381
preference, 1381
refuges,
Cotesia flavipes, 627
Diatraea saccharalis, 627
sentinel larvae, 627
sugarcane, 627
remote sensing, 1045, 1718
imported fire ants, 1718
mounds, 1045
multispectral, 1045
photointerpretation, 1718
repeatability,
costs of mating, 779
longevity, 779
sexual selection, 779
starvation resistance, 779
repellency,
catnip, 1562
elemol, 1562
nepetalactone, 1562
osage orange, 1562
resistance,
corn, 35
phenology, 35
rotation, 35
soybean, 35
resistance management, 681, 740, 930, 1102
alternate hosts, 681, 740, 1102
Bacillus thuringiensis, 1102
corn, 930
Diabrotica virgifera virgifera, 681, 740, 1102
elemental marking, 930
prairie grasses, 740
rootworm, 681
southwestern corn borer, 930
transgenic-rootworm maize, 1102
Zea mays, 930
resource availability,
dung beetles, 188
nesting, 188
ovarian condition, 188
sexual maturation, 188
resource partitioning,
foraging, 1602
polygyne, 1602
red imported fire ant, 1602
Solenopsis invicta, 1602
respiratory physiology,
isothermal calorimetry, 832
multicolored Asian lady beetle, 832
R_{CO2}, 832
temperature dependence, 832
Reticulitermes flavipes, 197, 1145
anaerobic protists, 1145
Coptotermes formosanus, 197
habitat temperature, 197
physiological saline solution, 1145
protist counting techniques, 1145
seasonal variation, 197
symbiotic protists, 1145
thermal tolerances, 197
Rhinoncomimus latipes,
feeding, 990
mile-a-minute weed, 990
Polygonum perfoliatum, 990
simulated herbivory, 990
Rhopalosiphum maidis,
Acyrthosiphon pisum, 1593
Aphis glycines, 1593
landscape ecology, 1593
Therioaphis trifolii, 1593
Rhyacionia frustrana,
damage prediction, 397
pheromone traps, 397
Pinus taeda, 397
risk,
natural habitats, 310
vineyards, 310
wild grape, 310
risk assessment, 958, 1109, 1116
biological control, 958
Bt corn, 1116
chronic effects, 1116
Danaus plexippus, 1109
monarch butterfly, 1116
nontarget, 1116
nontargets, 1109
pollinivory, 958
transgenic corn, 1109
transgenic crops, 958
Zea mays, 958
root feeding,
food conversion efficiency, 1169
invasive species, 1169
nutritional ecology, 1169
phytosterols, 1169
western corn rootworm, 1169
rootworm,
alternate hosts, 681
Diabrotica virgifera virgifera, 681
resistance management, 681
Rosa arkansana,
Diplolepis spp., 1052
plant-stress hypothesis, 1052
plant-vigor hypothesis, 1052
tall-grass prairie, 1052
rotation,
corn, 35
phenology, 35
resistance, 35
soybean, 35

S

S. molesta,
biological control, 587
Forelius, 587
Monomorium, 587
Pheidole, 587
Salix scouleriana,
Euura sp., 700
plant vigor, 700

population dynamics, 700
preference-performance linkage, 700
salt marsh,
caterpillar ecology, 1535
Coenonympha tullia nipisiquit, 1535
hypoxia, 1535
tidal submergence, 1535
sampling,
boll-feeding Lepidoptera, 1458
within-plant distribution, 1458
sampling techniques,
gypsy moth, 418
Lepidoptera, 418
population dynamics, 418
sawtoothed grain beetle,
olfaction, 75
Oryzaephilus surinamensis, 75
packaging, 75
stored products, 75
Scarabaeidae,
biological control, 520
host selection, 520
parasitoid, 520
turfgrass, 520
Scarabaeoidea,
assemblage composition, 45
dung beetle, 45
spatial variability, 45
trophic resource, 45
Schistocerca gregaria,
aggregation, 1577
gregarization, 1577
phase, 1577
Scolytidae,
Cytospora, 21
feltleaf willow, 21
Trypophloeus, 21
Scymnus loewii,
biological control, 282
conservation, 282
Hippodamia convergens, 282
seasonal adaptation,
bruchid, 1163
diapause, 1163
overwintering stage, 1163
seasonal development,
competition, 1232
diapause, 1232
Phoracantha recurva, 1232
Phoracantha semipunctata, 1232
seasonal variation,
Coptotermes formosanus, 197
habitat temperature, 197
Reticulitermes flavipes, 197
thermal tolerances, 197
semiochemicals,
invasive species, 206
monoterpenes, 206
phylogenetic geography, 206
sentinel larvae,
Cotesia flavipes, 627
Diatraea saccharalis, 627
refuges, 627
sugarcane, 627
sex ratio, 328, 1431
acoustic communication, 1431
host density, 328
host mortality, 328
mole cricket, 1431
moon phase, 1431
progeny production, 328
sexual maturation,
dung beetles, 188
nesting, 188
ovarian condition, 188
resource availability, 188
sexual selection, 657, 779
body size, 657
costs of mating, 779
longevity, 779
male aggression, 657
mating success, 657
Nadezhdiella cantori, 657
repeatability, 779
starvation resistance, 779
shrubs,
fertile islands, 599
microbes, 599
nitrogen, 599
spatial heterogeneity, 599
Silphium,
Asteraceae, 291
parasitoid, 291
prairie, 291
predation, 291
stable isotope, 291
similarity indexes,
Asteraceae, 297
floral visitors, 297
temporal variation, 297
xeric community, 297
simulated herbivory,
feeding, 990
mile-a-minute weed, 990
Polygonum perfoliatum, 990
Rhinoncomimus latipes, 990
Sipha elegans,
alate, 546
apterous, 546
yellow sugarcane aphid, 546
site-specific pest management,
spatial distribution, 1037
sunflower midge, 1037
Sitona hispidulus,
alfalfa, 119
clover root curculio, 119
Medicago sativa, 119
nodules, 119
Sitophilus,
Anisopteromalus calandrae, 671
Cephalonomia waterstoni, 671
Cryptolestes, 671
Pea protein, 671
Sitotroga cerealella,
corn, 807
life history, 807
modeling, 807
survivorship, 807
smoke tree sharpshooter,
diffusion model, 88
glassy-winged sharpshooter, 88
IgG protein markers, 88
yellow sticky traps, 88
sodium,
foraging, 1020
mud puddling, 1020
Pieris napi, 1020
soil moisture,
biological control, 1127
entomopathogenic fungi, 1127
Malacosoma disstria, 1127
temperature, 1127
soil properties,
Aethina tumida, 794
Apis mellifera, 794
pupation success, 794
soil survey,
Bacillus thuringiensis, 1335
Beauveria bassiana, 1335
Metarhizium anisopliae, 1335
Steinernema oregonense, 1335
soil water regimen,
Diptera, 1282
environmental stress, 1282
flooding, 1282
Isopoda, 1282
Solanum linnaeanum,
dispersal, 1722
metapopulation biology, 1722
oviposition preference, 1722
Solanum torvum, 1722
Solanum torvum,
dispersal, 1722
metapopulation biology, 1722
oviposition preference, 1722
Solanum linnaeanum, 1722
Solenopsis geminata,
biocontrol, 1436
host range, 1436
Solenopsis invicta, 1436
Solenopsis xyloni, 1436
Solenopsis invicta, 340, 975, 1436, 1602
biocontrol, 1436
biological control, 340
entomopathogen, 975
fire ant, 340
foraging, 1602
fungal pathogen, 975
host range, 1436
microbial control, 975
pathogen, 340
polygyne, 1602

polygyny, 340
red imported fire ant, 1602
resource partitioning, 1602
Solenopsis geminata, 1436
Solenopsis xyloni, 1436
Solenopsis molesta,
native ant, 1626
predation, 1626
red imported fire ant, 1626
thief ant, 1626
Solenopsis xyloni,
biocontrol, 1436
host range, 1436
Solenopsis geminata, 1436
Solenopsis invicta, 1436
South Korea,
Acer mono, 540
Anoplophora glabripennis, 540
Asian longhorned beetle, 540
invasive species, 540
southwestern corn borer, 487, 930
Bt maize, 487
corn, 930
dispersal, 487
elemental marking, 487, 930
resistance management, 930
Zea mays, 930
soybean,
corn, 35
phenology, 35
resistance, 35
rotation, 35
spatial autocorrelation,
Anoplophora glabripennis, 435
eradication, 435
invasive species, 435
mark–recapture, 435
spatial distribution, 1037, 1266
habitat distribution, 1266
Ixodes scapularis, 1266
site-specific pest management, 1037
sunflower midge, 1037
survival, 1266
spatial dynamics,
Aesculus hippocastanum, 1584
biological invasion, 1584
Cameraria ohridella, 1584
pheromone trapping, 1584
spatial heterogeneity,
fertile islands, 599
microbes, 599
nitrogen, 599
shrubs, 599
spatial processes,
forecasting, 257
geographic information system, 257
Markov chain, 257
pest management, 257
spatial variability,
assemblage composition, 45
dung beetle, 45
Scarabaeoidea, 45
trophic resource, 45
specialist,
adaptation, 1477
generalist, 1477
oviposition, 1477
polyphagy, 1477
species replacement,
fragmentation, 234
habitat loss, 234
matrix-effects, 234
Ostrinia nubilalis, 234
Plathypena scabra, 234
spectral quantum flux,
aphid-borne viruses, 1447
Aphis gossypii, 1447
Bemisia argentifolii, 1447
UV reflective mulch, 1447
wheat straw mulch, 1447
spinosad,
impatiens, 1344
inoculative augmentation, 1344
nematode rearing, 1344
western flower thrips, 1344
spiroplasma,
corn leafhopper, 1644
corn stunt, 1644
Dalbulus maidis, 1644
Spiroplasma kunkelii, 1644
Spiroplasma kunkelii,
corn leafhopper, 1644
corn stunt, 1644
Dalbulus maidis, 1644
spiroplasma, 1644
Spodoptera exigua, 54, 964
big-eyed bug, 964
entomopathogenic algae, 54
generalist predator, 964
Helicosporidium, 54
Helicoverpa zea, 964
Heliocoverpa zea, 54
life table, 964
Trichoplusia ni, 54
Spodoptera frugiperda,
fall armyworm, 881
host strains, 881
population dynamics, 881
squash leaf curl virus,
aphids, 1242
Bemisia tabaci, 1242
papaya ring spot virus, 1242
watermelon, 1242
stable isotope,
Asteraceae, 291
parasitoid, 291
prairie, 291
predation, 291
Silphium, 291
Staphylinidae,
biological control, 1681
Carabidae, 1681
turfgrass, 1681
voracity, 1681
starvation resistance,
costs of mating, 779
longevity, 779
repeatability, 779
sexual selection, 779
Steinernema oregonense,
Bacillus thuringiensis, 1335
Beauveria bassiana, 1335
Metarhizium anisopliae, 1335
soil survey, 1335
stink bugs,
boll rot, 1095
cotton, 1095
hard-locked, 1095
stopover,
migration, 249
monarchs, 249
timing, 249
weather, 249
stored grain,
Cryptolestes ferrugineus, 1003
distribution, 1003
movement, 1003
temperature gradient, 1003
wandering, 1003
stored products,
olfaction, 75
Oryzaephilus surinamensis, 75
packaging, 75
sawtoothed grain beetle, 75
stored-product beetles, 426, 1465
decision support tools, 426
modeling, 426
monitoring, 1465
Oklahoma, 426, 1465
trap catch analysis, 1465
straw mulch,
insects, 1632
potato, 1632
watermelon, 1632
weeds, 1632
subarctic,
phenological model, 1513
thermoregulation, 1513
sucrose,
Aphis gossypii, 690
cotton, 690
cotton aphid, 690
glucose, 690
leaf sugars, 690
sugarcane,
Cotesia flavipes, 627
Diatraea saccharalis, 627
refuges, 627
sentinel larvae, 627

summer generation,
cold hardiness, 113
fall generation, 113
Locusta migratoria, 113
postwinter population, 113
sunflower midge,
site-specific pest management, 1037
spatial distribution, 1037
supercooling point,
cold hardiness, 815
Harmonia axyridis, 815
low temperature, 815
overwintering, 815
superparasitism,
behavior, 362
external/internal marking, 362
host quality, 362
supplemental food sources,
conservation biological control, 619
Tiphia pygidialis, 619
Tiphia vernalis, 619
white grubs, 619
survival, 823, 1266
biological control, 823
cold storage, 823
habitat distribution, 1266
house fly, 823
Ixodes scapularis, 1266
pteromalids, 823
spatial distribution, 1266
survivorship,
corn, 807
life history, 807
modeling, 807
Sitotroga cerealella, 807
sweep-net sampling,
Coccinellidae, 535
visual sampling, 535
yellow sticky cards, 535
symbiosis,
bark beetles, 28
insect–fungal interaction, 28
Ophiostoma ips, 28
ophiostomatoid fungi, 28
symbiotic protists,
anaerobic protists, 1145
physiological saline solution, 1145
protist counting techniques, 1145
Reticulitermes flavipes, 1145
synovigenic,
Calpodes ethlius, 227
ectoparasitoid, 227
Elachertus scutellatus, 227
koinobiont, 227

T

tall-grass prairie,
Diplolepis spp., 1052
plant-stress hypothesis, 1052
plant-vigor hypothesis, 1052
Rosa arkansana, 1052
tarnished plant bug,
egg load, 1389
flight behavior, 1389
flight mill, 1389
vertical flight chamber, 1389
western tarnished plant bug, 1389
Temnochila chlorodia,
biological control, 1554
chemical ecology, 1554
Enoclerus lecontei, 1554
Enoclerus sphegeus, 1554
temperature, 799, 1127, 1137, 1155, 1549
biological control, 1127
calorimetry, 799
Chrysomela lapponica, 799
color morph, 799
Cydia (Grapholita) molesta, 1155
developmental duration, 1549
dispersal, 1155
entomopathogenic fungi, 1127
host plant, 1137
light phase, 1137
Lygus elisus, 1549
Macrolophus pygmaeus, 1137
Malacosoma disstria, 1127
photoperiod, 1155
population dynamics, 1549
predation rate, 1137
preimaginal environment, 1155
soil moisture, 1127
thermal balance, 799
temperature dependence,
isothermal calorimetry, 832
multicolored Asian lady beetle, 832
R_{CO2}, 832
respiratory physiology, 832
temperature gradient,
Cryptolestes ferrugineus, 1003
distribution, 1003
movement, 1003
stored grain, 1003
wandering, 1003
temperature-dependent development, 107, 787
egg hatch, 107
model, 107, 787
Ostrinia furnacalis, 787
oviposition behavior, 787
pea, 107
pea leaf weevil, 107
Trichogramma ostriniae, 787
temporal spatial dynamics,
alfalfa, 890
Empoasca fabae, 890
Medicago sativa, 890
proportion maps, 890
temporal variation,
Asteraceae, 297
floral visitors, 297
similarity indexes, 297
xeric community, 297
Tenodera aridifolia sinensis, 182, 1032
antipredator behavior, 1032
biological control, 1032
deme extinction, 182
introduced predator, 1032
introduced species, 182
mantids, 182
metapopulations, 182
Pardosa milvina, 1032
termite baiting system,
Formosan subterranean termite, 242
monitoring station, 242
native subterranean termite, 242
termite foraging activity, 242
termite foraging activity,
Formosan subterranean termite, 242
monitoring station, 242
native subterranean termite, 242
termite baiting system, 242
Tetragnatha,
Cyclosa, 1296
edge effects, 1296
Hawaii, 1296
lava flows, 1296
Tetropium fuscum,
attractants, 844
Cerambycidae, 844
host volatiles, 844
traps, 844
Thamnurgus euphorbiae,
beetle, 1673
biological control, 1673
Coleoptera, 1673
weed, 1673
The Netherlands,
Ageniaspis fuscicollis, 528
field sex ratio, 528
Yponomeuta, 528
Therioaphis trifolii,
Acyrthosiphon pisum, 1593
Aphis glycines, 1593
landscape ecology, 1593
Rhopalosiphum maidis, 1593
thermal balance,
calorimetry, 799
Chrysomela lapponica, 799
color morph, 799

temperature, 799
thermal tolerances,
Coptotermes formosanus, 197
habitat temperature, 197
Reticulitermes flavipes, 197
seasonal variation, 197
thermoregulation,
phenological model, 1513
subarctic, 1513
thief ant,
native ant, 1626
predation, 1626
red imported fire ant, 1626
Solenopsis molesta, 1626
threshold,
development, 1
model, 1
Nephus bisignatus, 1
Nephus includens, 1
thrips,
hoverflies, 1416
light-emitting diodes, 1416
traps, 1416
tidal submergence,
caterpillar ecology, 1535
Coenonympha tullia nipisiquit, 1535
hypoxia, 1535
salt marsh, 1535
timing,
migration, 249
monarchs, 249
stopover, 249
weather, 249
Tiphia pygidialis,
conservation biological control, 619
supplemental food sources, 619
Tiphia vernalis, 619
white grubs, 619
Tiphia vernalis,
conservation biological control, 619
supplemental food sources, 619
Tiphia pygidialis, 619
white grubs, 619
Tomicus piniperda,
disruptants, 221
nonhost volatiles, 221
verbenone, 221
Tortricidae, 506, 513
Adoxophyes orana, 513
diapause termination, 513
Grapholita prunivora, 506
lesser apple worm, 506
overwintering, 513
photoperiod, 513
toxic effects,
amorphogenic effects, 1151
Apis mellifera, 1151
fungicides, 1151
honey bee, 1151
transgenic corn,
Danaus plexippus, 1109
nontargets, 1109
risk assessment, 1109
transgenic crop,
Bacillus thuringiensis, 1025
diamondback moth, 1025
insect behavior, 1025
Plutella xylostella, 1025
transgenic crops,
biological control, 958
pollinivory, 958
risk assessment, 958
Zea mays, 958
transgenic plants,
Bacillus thuringiensis, 982
corn, 982
fall armyworm, 982
nucleopolyhedrovirus, 982
transgenic-rootworm maize,
alternate hosts, 1102
Bacillus thuringiensis, 1102
Diabrotica virgifera virgifera, 1102
resistance management, 1102
trap catch analysis,
monitoring, 1465
Oklahoma, 1465
stored-product beetles, 1465
trap crop,
corn earworm, 1371
cotton, 1371
grain sorghum, 1371
Helicoverpa zea, 1371
traps, 844, 1416
attractants, 844
Cerambycidae, 844
host volatiles, 844
hoverflies, 1416
light-emitting diodes, 1416
Tetropium fuscum, 844
thrips, 1416
Trichogramma,
egg parasitoids, 855
premating dispersion, 855
virgin females, 855
Trichogramma ostriniae,
model, 787
Ostrinia furnacalis, 787
oviposition behavior, 787
temperature-dependent development, 787
Trichoplusia ni,
entomopathogenic algae, 54
Helicosporidium, 54
Heliocoverpa zea, 54
Spodoptera exigua, 54
trophic preferences,
cattle dung, 370
dung beetles, 370
horse dung, 370
olfactometer, 370
trophic resource,
assemblage composition, 45
dung beetle, 45
Scarabaeoidea, 45
spatial variability, 45
Trypophloeus,
Cytospora, 21
feltleaf willow, 21
Scolytidae, 21
turf,
conservation biological control, 1662
habitat manipulation, 1662
turfgrass, 520, 1681
biological control, 520, 1681
Carabidae, 1681
host selection, 520
parasitoid, 520
Scarabaeidae, 520
Staphylinidae, 1681
voracity, 1681
twospotted spider mite,
aphids, 1014
Coleomegilla, 1014
Geocoris, 1014
Orius, 1014
predators, 1014

U

UV irradiation,
Evania appendigaster, 1321
freezing, 1321
gamma irradiation, 1321
heating, 1321
Periplaneta americana, 1321
UV reflective mulch,
aphid-borne viruses, 1447
Aphis gossypii, 1447
Bemisia argentifolii, 1447
spectral quantum flux, 1447
wheat straw mulch, 1447

V

Vaccinium,
Grapholita, 405
lingonberry fruitworm, 405
pheromone traps, 405
population monitoring, 405
Vaccinium ashei,
Apis mellifera, 100
nectar robbing, 100
rabbiteye blueberry, 100
Xylocopa virginica, 100
vegetable,
Bemisia tabaci, 839
biological control, 839
Delphastus catalinae, 839
whitefly, 839

vegetative diversity,
European corn borer, 320
natural enemies, 320
Ostrinia nubilalis, 320
weeds, 320
verbenone,
disruptants, 221
nonhost volatiles, 221
Tomicus piniperda, 221
vertical flight chamber,
egg load, 1389
flight behavior, 1389
flight mill, 1389
tarnished plant bug, 1389
western tarnished plant bug, 1389
Vigna unguiculata,
canopy temperature, 471
defoliation, 471
pod position, 471
vineyards,
natural habitats, 310
risk, 310
wild grape, 310
virgin females,
egg parasitoids, 855
premating dispersion, 855
Trichogramma, 855
virus vectors,
coat protein–mediated resistance, 1207
host plant resistance, 1207
insect behavior, 1207
virus-induced volatiles, 1207
virus-induced volatiles,
coat protein–mediated resistance, 1207
host plant resistance, 1207
insect behavior, 1207
virus vectors, 1207
vision, 81, 1410
Anthonomus pomorum, 1410
chromatic contrast, 81
colors, 1410
host finding, 1410
host location, 81
host plant, 1410
pupal parasitoid, 81
wavelength discrimination, 81
visual sampling,
Coccinellidae, 535
sweep-net sampling, 535
yellow sticky cards, 535
voracity,
biological control, 1681
Carabidae, 1681
Staphylinidae, 1681
turfgrass, 1681

W

wandering,
Cryptolestes ferrugineus, 1003
distribution, 1003
movement, 1003
stored grain, 1003
temperature gradient, 1003
watermelon, 1242, 1632
aphids, 1242
Bemisia tabaci, 1242
insects, 1632
papaya ring spot virus, 1242
potato, 1632
squash leaf curl virus, 1242
straw mulch, 1632
weeds, 1632
watermilfoil,
biocontrol, 872
development time, 872
Euhrychiopsis lecontei, 872
wavelength discrimination,
chromatic contrast, 81
host location, 81
pupal parasitoid, 81
vision, 81
weather,
migration, 249
monarchs, 249
stopover, 249
timing, 249
webspinners,
detritivore, 907
insect ecology, 907
ordination, 907
weed,
beetle, 1673
biological control, 1673
Coleoptera, 1673
Thamnurgus euphorbiae, 1673
weed biological control,
Christmas berry, 1351
herbivory, 1351
leafroller, 1351
life history, 1351
weeds, 320, 577, 1632
European corn borer, 320
field corn, 577
Gryllus spp., 577
Harpalus pensylvanicus, 577
insects, 1632
Lycosidae, 577
natural enemies, 320
Ostrinia nubilalis, 320
potato, 1632
straw mulch, 1632
vegetative diversity, 320
watermelon, 1632
western corn rootworm, 650, 1169
Diabrotica virgifera virgifera, 650
diel periodicity, 650
dispersal, 650
flight, 650
food conversion efficiency, 1169
invasive species, 1169
nutritional ecology, 1169
phytosterols, 1169
root feeding, 1169
western flower thrips,
impatiens, 1344
inoculative augmentation, 1344
nematode rearing, 1344
spinosad, 1344
western tarnished plant bug,
egg load, 1389
flight behavior, 1389
flight mill, 1389
tarnished plant bug, 1389
vertical flight chamber, 1389
wheat straw mulch,
aphid-borne viruses, 1447
Aphis gossypii, 1447
Bemisia argentifolii, 1447
spectral quantum flux, 1447
UV reflective mulch, 1447
white grubs,
conservation biological control, 619
supplemental food sources, 619
Tiphia pygidialis, 619
Tiphia vernalis, 619
whitefly,
Bemisia tabaci, 839
biological control, 839
Delphastus catalinae, 839
vegetable, 839
wild grape,
natural habitats, 310
risk, 310
vineyards, 310
wing loading,
dispersal-reproduction syndrome, 1471
energy reserves, 1471
host plant quality, 1471
Phthorimaea operculella, 1471
within-plant distribution,
boll-feeding Lepidoptera, 1458
sampling, 1458

X

xeric community,
Asteraceae, 297
floral visitors, 297
similarity indexes, 297
temporal variation, 297
Xylocopa virginica,
Apis mellifera, 100
nectar robbing, 100
rabbiteye blueberry, 100

Vaccinium ashei, 100

Y

yellow sticky cards,
Coccinellidae, 535
sweep-net sampling, 535
visual sampling, 535
yellow sticky traps,
diffusion model, 88
glassy-winged sharpshooter, 88
IgG protein markers, 88
smoke tree sharpshooter, 88
yellow sugarcane aphid,
alate, 546
apterous, 546
Sipha elegans, 546
Yponomeuta,
Ageniaspis fuscicollis, 528
field sex ratio, 528
The Netherlands, 528

Z

Zea mays, 930, 958, 1192
biological control, 958
corn, 930
elemental marking, 930
insect vector, 1192
morphological variation, 1192
pollinivory, 958
resistance management, 930
risk assessment, 958
southwestern corn borer, 930
transgenic crops, 958